

# Cardiovascular events or unexpected death with 60%-70%

C

[Business](#), [Management](#)



Cardiovascular disease constitutes the frequent dominant cause of worldwide death affecting the youthful population (1) On clinical entities silent MI annually experience CHD with the approximate of 9.

8 million per year predicting the unrecognized symptoms related to the phenomenon of either ambulatory ischemic events or unexpected death with 60%-70% asymptomatic manifestation. (2) The kidney disease prediction on clinical practice guidelines proves the demonstration of ischemic chest pain is a strong venue of cardiovascular morbidity measures and all cause-mortality (3) with the classification of hemodynamic moderations independently calculate glomerular filtration rate (eGFR) on the scheme of arbitrary renal stages altering albuminuria, hyperfiltration and hyperglycemia in renal failure. Consequently CKD as a clinical syndrome equivalent to CHD investigate the urinary excretion proteins and TGF-beta 1 initially in nephropathy as a diagnostic value in clinical parameters of previous MI in diabetic individuals. (4) In the Framingham heart study, the estimation of silent ischemia detection on large scale misinterpreted by atypical angina following normal ECG with the unnoticeable prevalence in broad ranging population <45 years young men and women including the illness of metabolic impairments, eGFR fall, advanced CKD at dialysis, chronic inflammations and prothrombic attributes in the acceptable autopsy reports. (5) Additionally diabetic kidney disease progress with the albuminuria status and serum creatinine measurements on screening at initial conventional method in pubertal diabetic interval challenging hyperglycemia on clinical nephropathy. (6) According to the Epidemiological controversies, poor estimation of CAD in CKD cases at less ratios make the diagnostic differences

by the interpretation of ECG changes, risk factor profiles, pain perception, cardiac biomarkers and MI perfusions assessment in the objective of angina culminate the presentation of constant changes in troponin values represent the attributes of premature CHD by the conclusion of progressive atheromatous plaques and calcifications on the likelihood of sepsis, anemia, platelet aggressiveness, nitric oxide abnormal metabolism, arterial stiffness, calcium-phosphate homeostasis, endothelial dysfunctions, recreational drugs, history of multiple traumas and surgeries.

Classically, we present the case of inferior MI in young women with type 2 diabetes evolutions of 10 years and previous heart failure cause to death prolonging the complications of dyslipidemia progressive to the advancement of diabetic nephropathy in extending membranoproliferative glomerulonephritis. Case presentation A 35-year-old female present to Emergency Department with severe chest pain, palpitation and vomiting for 3 days. She has been diagnosed with the previous episodes of heart failure and traditional risk factors of CAD in medical history. She described her chest pain with tightness and flank dull pain at both the areas of kidneys with back pain.

On physical examination, Cardiac sounds were normal on auscultation with no tenderness on palpation, no intra-abdominal rebound masses, no neck stiffness, no jugular vein enlargement, no dysmenorrhea, no clubbing, no family history of CAD and no hypertension. She was profound sweating on presence with weight loss, urine retention, fatigability and restlessness from 1 week. Her medications at the time of review include Aspirin, Statin,

Metformin, Insulin, Diuretics and Omeprazole. At Admission, BP was 85/60mmHg and her heartrates 66 bpm.

ECG showed normal sinus rhythm with ST elevation in leads II, III and avf with the reciprocal of ST segment depression in leads V1-V6 as shown in Fig 1A

The crucial step for ruling out myocardial injury, clinical diagnosis begin with the measurement of cardiac enzymes as shown in Table 1 Moreover on the primary assessment of troponin elevation and NT-proBNP impairment assess the specificity and sensitivity limitations on trans-thoracic

echocardiography revealing hypokinesia with an LVEF of 48% motion index on homogenous contrast reflecting MI tension on inferior walls suspecting

intracardiac thrombosis and pulmonary embolism. Figure 1 (A) Initially ECG shows ST elevation at inferior leads with the reciprocal of ST depression in avR. (B) No simultaneous changes in right ventricular MI on various

segments of ECG. (C) New ST depression in the leads of II, III, and

avF after following fibrinolysis in 12 lead ECG. Table 1 Clinical values of

Combined Detection of 5 Indicators in the Diagnosis of Acute MI.

In the suspicion of fibrinolysis and thrombo embolism, thoracic

ultrasonography TUS certainly performed prior to the normal chest imaging previously and false positive predictive value in D-dimer test as shown in

Table 2 On the emergency based history of angina, bilateral thoracic

probe examine the presence of the left sided non specific pleural lesion of more than 5mm on screening. It provokes the follow up of thrombolytic with

the association of hypotension. Therefore, anticoagulation includes low-molecular-weight heparin therapy (LMWH) and tPA produce successful

reperfusion within 12hrs non-invasively. Table 2 Quantitative D-dimer Assay for Pulmonary Embolism Diagnostic Test. In regards with Gastrointestinal aspects, the alarming signs of dehydration, nausea, vomiting, fatigue and back pain warrant the examination of a comprehensive metabolic panel and amylase, lipase testing for the consideration of gastroenteritis or acute pancreatitis. The normal values result self-limiting bacterial infections by the management of fluid replacement, Calcitonin and supportive care.

As the patient on type 2 diabetes expansion on clinical estimation follow urinalysis on palpation of the bladder and oliguria. According to the quantitative measurements on total protein positive test, the exercising ECG reviewed on high standards verify the reciprocal changes in pathologic Q waves and hyperacute T waves in nonfatal angina attack reflect preload independently as shown in figure 1B. Apart from the renal profile, further globulin tests were progressed on the basis of laboratory evidences as shown in Table 3 - 4 decline in eGFR, leucocytosis and elevated cholesterol conclude the pathogenesis of contrast induced nephropathy in association of nephrotoxic drugs eliminating the advanced staging of kidney damage other than glomerulonephritis and residual renal dysfunctions. Table 3 Comprehensive Metabolic Panel with eGFR Blood Test. Table 4 Complete Blood Count Test Results. On the basis of ANA negative investigation, monoclonal immunoglobulin IgG determine the pre-malignancy in renal insufficiency with plasmapheresis at high risk of multiple myelomas as shown in Table 5 here in the diagnosis of proteinuria and myeloma related diseases Bence Jones test reveal false negative results in concentrated urine.

At result, vitamin K status in CKD sub clinically link to the formation of arterial calcification in the highmoderations of atherosclerosis constitute the notable limitations on independent peritoneal dialysis to maintain the equivalent nutrition at the less co-morbidity of young age in CKD. Table 5 Serum Protein Electrophoresis to diagnose M protein. Differential Diagnosis Prinzmetals angina/vasospasm, cardiogenic shock, cardiac contusion, pulmonary edema, acute gastritis, GERD and anxiety disorders are unlikely considered on pursued clinical presentation as reviewed. Treatment Management is initiative with the long-lasting insulin therapy in type 2 diabetes with the combination of Sulfonylurea and Metformin to control hyperglycemia. Secondly use of diuretics to restore electrolyte imbalance and Vitamin C for the nausea feeling. Thirdly Diazepam orally for the anxiety and cardiac therapy Cedilanid for hemodynamic stability, Dopamine hydrochloride for improving the cardiac functions, Hydroxylamine and MgSO<sub>4</sub> to control frequent arrhythmias, Clopidogrel 150mg + Aspirin 100mg with heparin therapy of LMWH in the preventions of heart failure and recurrent myocardial infarction. Lastly IV Sodium bicarbonate + insulin + 50% Dextrose for hyperkalemia and Atorvastatin of 20mg oral/day for LDL reduction.

Follow up On the Ninth day, ECG changes as shown in fig 1C, ST resolution and T wave inversion after the pharmaceutical drugs. At practical measures IV human albumin infusion as a therapeutic plasmapheresis remarkably improved the tailored indication of hypovolemic shock in the significance of cardiac improvement. Hence at the objective of primary care with proper monitoring of stable renal functions by calcium gluconate, on fifteenth day

patient discharged with effective diet planning assumed by community-based clinicians in providing selfmanagement to control delicate balance in postprandial hyperglycemia adjustments.

Discussion The DIAD (Ischemia detection in asymptomatic Diabetics) (5, 7) assumes the importance of greater incidence in long standing type 2 diabetes mellitus focus on the factor of occlusion in arteries on the possibility of judicious analyses with no support of scientific data in the management of anti-ischemic medications at frequent CAD cases. Hence, the investigations of massive consequences intermediate undoubtedly on clinical scoring as addressed for the issue of positive prognostic screening program in the upcoming studies. American Diabetes Association (ADA) recommend the measures of Beta blockers or re-vascularization medical therapy on aggressive intensive treated cases on investigating the annual review of abnormal resting ECG with the lesser degrees of ischemia intervention can improve the prognosis on cardiovascular events.

The nontraditional factors of hyper coagulation and clotting mediators (8) pronounce the elevation of high risk on thrombotic events statistically with the complications of CKD underlying the unclear etiology predominantly result congestive heart failure, ESRD, hemorrhagic stroke and relative risk of peripheral artery disease proportional to sudden cardiac death. Thus, an appropriate medical therapeutic management needed in terms of risk factors incidental preventions in adult onset diabetes. (9) In primary prevention study at Helsinki Heart Study (10) show poor outcomes in Diabetic individuals with CHD identifying high risk of aggressiveness in dyslipidemia treatment for

the maintenance of LDL and Total protein target the statin drugs as a pharmacological intervention for the trials as a first choice in young diabetic nephropathy patients. The General Practice Research thrombosis Prevention trial (11) on the secondary prevention confirm the benefit of Aspirin treatment in the establishment of atherosclerotic disease in prospective trials reduced the risk of CHD and non fatal events on the clinical recommendation of anti-platelet therapy can also be used as a preventive strategy to overt the nephropathy in <30 years age individuals. Therefore large phase prospective studies and trials are required to explain the issues of uncertain protein restriction in the adherence of management in routine setting care in diabetic nephropathy. Observational Studies in the demonstration of direct effect on CVD risk factors deteriorate the kidney functions in hyperglycemia.

The Reduction of End points in Non-insulin-dependent Diabetes with the Angiotensin II Antagonist Losartan (RENAAL) and Irbesartan Diabetic Nephropathy Trial (IDNT) studies include the trial of Losartan and Irbesartan as a renoprotective in the combination of Ramipril and Telmisartan initiate the defensive effect on proteinuria as compared to the therapy of (VA NEPHRON-D) study of Losartan and Lisinopril on macroalbuminuria > 300mg/day. Thus, the supportive directions on definite limitations of safety concerns utilize the consideration of Renin Angiotensin Aldosterone System (RAAS) lessens microalbuminuria 30-300mg/day in normid diabetes cases. (12)

Hyperglycemia as a therapeutic potent in diabetes, the epidemiological early analysis illustrate the fundamental controversy of minimal outcomes in macrovascular hazards can ascend the occasion of CVD risk factors,



extravagant mortality rates and vigorous symptoms with the median of HbA1c%. The Action in Controlling Cardiac Risk factors in Diabetes (ACCORD) present the current affirmation of delaying vascular complications related to the consequences of CKD staging 3-4 can be patently achieved by the optimal goal of HbA1c and hypoglycemia incidents. Accordingly, a tight control on hyperglycemia is permeable to convert the high risk of hyperfiltration and glomerular hypertrophy partially on HbA1c <7% and apparent supremacy to control the normal ranges with the treatment of insulin in the maintenance of proteinuria on reduced value. According to the American Heart Association guidelines, the pharmacotherapy in CKD associated with CVD risk factors include the counsel use of Fibrinolytic, Antiplatelet, Glycoprotein II b/III a receptor antagonist, Anti-coagulants, Beta blockers, ACEIs/ARBs, Aldosterone blocker and Statin can assess the randomized controlled trials of efficacy and welfare to diminish the vascular events in non chronic dialysis patients.

The another Study of Heart and Renal Protection (SHARP) involve the substantial results in combined therapy composite to the dominance in controlling the major atherosclerotic relative risks, intracranial hemorrhage, left ventricular hypertrophy and STEMI intimated the remarkable decline in hospital death and sudden cardiac arrest for least 1 year. Ultimately, pharmacokinetic studies in renal dysfunction require essential regulations for the clinical controlled trials further on extensive population with distinct precise dosing in terminating the predictable adverse outcome pathways.