

Deep brain stimulation for movement disorders – a history of success and challeng...

[Health & Medicine](#)



**ASSIGN
BUSTER**

A commentary on

Fasano, A., Herzog, J., Raethjen, J., Rose, F. E. M., Muthuraman, M., Volkman, J., Falk, D., Elble, R., and Deuschl, G. (2010a). Gait ataxia in essential tremor is differentially modulated by thalamic stimulation. *Brain* 133, 3635–3648.

Fasano, A., Romito, L. M., Daniele, A., Piano, C., Zinno, M., Bentivoglio, A. R., and Albanese, A. (2010b). Motor and cognitive outcome in patients with Parkinson's disease 8 years after subthalamic implants. *Brain* 133, 2664–2676.

Gradinaru, V., Mogri, M., Thompson, K. R., Henderson, J. M., and Deisseroth, K. (2009). Optical deconstruction of parkinsonian neural circuitry. *Science* 324, 354–359.

Okun, M. S., Gallo, B. V., Mandybur, G., Jagid, J., Foote, K. D., Revilla, F. J., Alterman, R., Jankovic, J., Simpson, R., Junn, F., Verhagen, L., Arle, J. E., Ford, B., Goodman, R. R., Stewart, R. M., Horn, S., Baltuch, G. H., Kopell, B. H., Marshall, F., Peichel, D., Pahwa, R., Lyons, K. E., Tröster, A. I., Vitek, J. L., Tagliati, M., and for the SJM DBS Study Group. (2012). Subthalamic deep brain stimulation with a constant-current device in Parkinson's disease: an open-label randomised controlled trial. *Lancet Neurol.* 11, 140–149.

Pizzolato, G., and Mandat, T. (2012). Deep brain stimulation for movement disorders. *Front. Integr. Neurosci.* 6: 2. doi: 10. 3389/fnint. 2012. 00002

Rouse, A. G., Stanslaski, S. R., Cong, P., Jensen, R. M., Afshar, P., Ullestad, D., Gupta, R., Molnar, G. F., Moran, D. W., and Denison, T. J. (2011). A chronic <https://assignbuster.com/deep-brain-stimulation-for-movement-disorders-a-history-of-success-and-challenges-to-conquer/>

generalized bi-directional brain-machine interface. *J. Neural Eng.* 8, 036018.
doi: 10.1088/1741-2560/8/3/036018

Sarem-Aslani, A., and Mullett, K. (2011). Industrial perspective on deep brain stimulation: history, current state, and future developments. *Front. Integr. Neurosci.* 5: 46. doi: 10.3389/fnint.2011.00046

Weaver, F. M., Follett, K., Stern, M., Hur, K., Harris, C., Marks, W. J., Rothlind, J., Sagher, O., Reda, D., Moy, C. S., Pahwa, R., Burchiel, K., Hogarth, P., Lai, E. C., Duda, J. E., Holloway, K., Samii, A., Horn, S., Bronstein, J., Stoner, G., Heemskerk, J., Huang, G. D., and CSP 468 Study Group. (2009). Bilateral deep brain stimulation vs best medical therapy for patients with advanced Parkinson disease: a randomized controlled trial. *JAMA* 301, 63–73.

Williams, A., Gill, S., Varma, T., Jenkinson, C., Quinn, N., Mitchell, R., Scott, R., Ives, N., Rick, C., Daniels, J., Patel, S., Wheatley, K., and PD SURG Collaborative Group. (2010). Deep brain stimulation plus best medical therapy versus best medical therapy alone for advanced Parkinson's disease (PD SURG trial): a randomised, open-label trial. *Lancet Neurol.* 9, 581–591.

Wojtecki, L., Vesper, J., and Schnitzler, A. (2011). Interleaving programming of subthalamic deep brain stimulation to reduce side effects with good motor outcome in a patient with Parkinson's disease. *Parkinsonism Relat. Disord.* 17, 293–294.