

Commentary on genetic mechanisms of antimicrobial resistance in bacteria from u.s...

[Health & Medicine](#)



**ASSIGN
BUSTER**

A commentary on

Genetic mechanisms of antimicrobial resistance identified in *Salmonella enterica*, *Escherichia coli*, and *Enterococcus* spp. isolated from U. S. food animals

by Frye, J. G., and Jackson, C. R. (2013). *Front. Microbiol.* 4: 135. doi: 10.3389/fmicb. 2013. 00135

While Frye and Jackson do a good job describing the common mechanisms of resistance found in food animals in the U. S., they err in stating that in the U. S. extended spectrum β -lactamases (ESBLs) “ thus far have only been found in human and not food animal isolates.”

In fact, Wittum et al., in 2010 first reported the collection of *Escherichia coli* expressing CTX-M-type ESBLs in fecal isolates from healthy cattle at a livestock market and from a diagnostic isolate submitted to the Ohio Animal Disease Diagnostic Laboratory. Since then, CTX-M-type ESBLs have been found in *E. coli* collected from healthy dairy calves in the western U. S. ([Davis et al., 2011](#)) and from 5 of 20 dairy farms in Ohio ([Mollenkopf et al., 2012](#)); in clinical *Salmonella enterica* isolates from swine in Minnesota and from turkeys in 4 states ([Wittum et al., 2012](#)); in *E. coli* from swine finishing barns in Michigan and Ohio and in *Klebsiella pneumoniae* from swine in Illinois ([Mollenkopf et al., 2013](#)).

We are not aware of studies finding CTX-M-type ESBLs in isolates from broiler chickens, but a retail chicken meat *E. coli* isolate expressing CTX-M-type ESBLs from Pennsylvania has been reported ([Doi et al., 2010](#)). Given the situation in other livestock species and in other countries, we expect the lack

<https://assignbuster.com/commentary-on-genetic-mechanisms-of-antimicrobial-resistance-in-bacteria-from-us-food-animals-esbls-are-here/>

of detection of CTX-M-type ESBLs in U. S. chicken isolates is more the result of lack of appropriate studies than a real absence of these on chicken farms. In addition to CTX-M-type ESBLs, the study of dairy calves in the western U. S. also found *E. coli* expressing OXA-type ESBLs ([Davis et al., 2011](#)).

References

Davis, M. A., Besser, T. E., Orfe, L. H., Baker, K. N., Lanier, A. S., Broschat, S. L., et al. (2011). Genotypic-phenotypic discrepancies between antibiotic resistance characteristics of *Escherichia coli* isolates from calves in management settings with high and low antibiotic use. *Appl. Environ. Microbiol.* *77*, 3293–3299. doi: 10.1128/AEM.02588-10

Doi, Y., Paterson, D. L., Egea, P., Pascual, A., López-Cerero, L., Navarro, M. D., et al. (2010). Extended-spectrum and CMY-type beta-lactamase-producing *Escherichia coli* in clinical samples and retail meat from Pittsburgh, USA and Seville, Spain. *Clin. Microbiol. Infect.* *16*, 33–38. doi: 10.1111/j.1469-0691.2009.03001.x

Mollenkopf, D. F., Mirecki, J. M., Daniels, J. B., Funk, J. A., Henry, S. C., Hansen, G. E., et al. (2013). *Escherichia coli* and *Klebsiella pneumoniae* producing CTX-M cephalosporinase from swine finishing barns and their association with antimicrobial use. *Appl. Environ. Microbiol.* *79*, 1052–1054. doi: 10.1128/AEM.03169-12

Mollenkopf, D. F., Weeman, M. F., Daniels, J. B., Abley, M. J., Mathews, J. L., Gebreyes, W. A., et al. (2012). Variable within- and between-herd diversity of

CTX-M cephalosporinase-bearing *Escherichia coli* isolates from dairy cattle.

Appl. Environ. Microbiol. 78, 4552–4560. doi: 10.1128/AEM.00373-12

Wittum, T. E., Mollenkopf, D. F., and Erdman, M. M. (2012). Detection of *Salmonella enterica* isolates producing CTX-M Cephalosporinase in U. S. livestock populations. *Appl. Environ. Microbiol.* 78, 7487–7491. doi: 10.1128/AEM.01682-12

Wittum, T. E., Mollenkopf, D. F., Daniels, J. B., Parkinson, A. E., Mathews, J. L., Fry, P. R., et al. (2010). CTX-M-type extended-spectrum β -lactamases present in *Escherichia coli* from the feces of cattle in Ohio, United States. *Foodborne Pathog. Dis.* 7, 1575–1579. doi: 10.1089/fpd.2010.0615