

Physical activity, sports practice, and cognitive functioning: the current resear...

[Health & Medicine](#)



**ASSIGN
BUSTER**

Physical Activity, Sports, and the Brain

The relationship between physical exercise and cognitive functioning has received much attention in recent years ([Northey et al., 2018](#) ; [Moran et al., 2019](#)). This has been an object of interest for decades, but much remains unknown. Neuroscience has advanced significantly, improving knowledge of brain functioning in response to different situations and its evolution over the course of people's lives ([Cabeza et al., 2018](#) ; [Dumoulin et al., 2018](#)). Scientists studying physical activity and sports have integrated this knowledge of brain functioning, using it to explain the contribution of physical exercise and how cognitive performance may increase performance in certain facets of sport ([Fink et al., 2018](#) ; [Hsu et al., 2018](#)).

Techniques such as electroencephalography ([Cheron et al., 2016](#) ; [Gutmann et al., 2018](#)), functional magnetic resonance imaging ([Chaddock-Heyman et al., 2013](#) ; [Fontes et al., 2015](#) ; [Chen et al., 2016](#)), positron emission tomography ([Boecker and Drzezga, 2016](#)), single photon emission tomography ([Shih et al., 2019](#)), or magnetoencephalography ([Huang et al., 2016](#)) have all improved the visualization and understanding of cognitive processes generated and developed in physical activity and sport contexts. The core of their contribution is the observation of brain changes during exercise, the impact of various tasks, and improvements in physical condition, which reflects this phenomenon ([Becker et al., 2016](#) ; [Jonasson et al., 2017](#) ; [Schwarb et al., 2017](#)).

In addition, research based on these techniques is complemented by other procedures developed in neurobiology and neurophysiology to explain

changes in the brain that are attributable to physical exercise. Specifically, hypotheses such as neurogenesis, synaptogenesis, or angiogenesis and the action of biomolecules such as irisin, brain-derived neurotrophic factor (BDNF), insulin-like growth factor-1 (IGF-1), vascular endothelial growth factor (VEGF), cathepsin B or interleukin-6 ([Tari et al., 2019](#)) have been proposed. However, these theoretical foundations are still in the process of consolidation for humans. In addition, some of these hypotheses arouse controversy among researchers, such as neurogenesis in humans throughout the life cycle, though this should be resolved in the coming years ([Sorrells et al., 2018](#) ; [Lourenco et al., 2019](#) ; [Pontifex et al., 2019](#) ; [Voss et al., 2019](#)).

In addition to an increased number of techniques available to assess the connection between physical exercise and brain functioning, other factors may have influenced the proliferation of this line of research. On the one hand, there have been warnings about increased sedentariness of lifestyles in some societies, prompting recommendations for physical activity to improve health among various sections of the population ([Hynynen et al., 2016](#) ; [Zylke and Bauchner, 2016](#)). In this regard, it has been proposed that brain functioning would benefit from physical activity, allowing better development in childhood and adolescence or acting as a protector in aging processes ([Erickson et al., 2015](#) ; [Costigan et al., 2016](#)). On the other hand, elite sport requires increased performance from athletes, thus encouraging the search for variables to improve the probability of success in competition. Cognitive functioning therefore has been an area of knowledge to which numerous researchers have aimed to contribute ([Krenn et al., 2018](#) ; [Sakamoto et al., 2018](#)).

In professional sports, differences between athletes are sometimes very subtle. Although it is difficult to eliminate all uncertainty from their performance, technical staff and research groups have sought to analyze and identify variables influencing this outcome. Thus, it is unsurprising that there are innumerable studies of physical preparation, technical and tactical aspects, or psychological influences ([Fister et al., 2018](#) ; [Dalen et al., 2019](#) ; [Henriksen et al., 2019](#)). Among these studies, brain analysis has attracted intense attention in recent years, becoming a prolific field of study and application that will probably continue in future as technical resources are perfected.

Physical Activity and Cognitive Functioning in Children, Adolescents, and the Elderly

Childhood and Adolescence

Analysis of the benefits of physical activity on people is especially relevant in childhood and adolescence. These are stages when future lifestyles are established, and acquired habits have a very strong influence on the state of health throughout life. However, during childhood and adolescence the brain is under construction and requires appropriate learning processes. In addition, children and adolescents are in a phase when their personal and social development is conditioned by multiple changes, to which they must make efforts to adapt. Therefore, the possible benefits of the cerebral functioning that physical exercise could produce become essential elements for children's and adolescents' growth and integration into the environment ([Wenner et al., 2013](#) ; [Lubans et al., 2016](#)). In fact, adequate cognitive

development during early stages is thought to contribute to improvements in wellbeing and mental health in adulthood ([Gale et al., 2012](#)).

In recent years, multiple studies have highlighted significant associations between physical practice and abilities, such as attention and concentration, executive functions, cognitive functioning speed, memory, or language (e. g., [Chaddock et al., 2011](#) ; [Scudder et al., 2014](#) ; [Donnelly et al., 2016](#) ; [Li et al., 2017](#) ; [Xue et al., 2019](#)). Various investigations have analyzed the acute effects of physical exercise ([Hillman et al., 2009b](#) ; [Elleberg and St-Louis-Deschênes, 2010](#) ; [Chang et al., 2012a](#)), the effects of a prolonged exercise program ([Hillman et al., 2014](#) ; [Reloba-Martínez et al., 2017](#)), as well as correlations between cognitive functioning with people engaged in regular physical activity or physical fitness exercises ([Hillman et al., 2005](#) ; [Carson et al., 2016](#) ; [Pérez-Lobato et al., 2016](#)).

Numerous papers have studied the relationship between physical activity and cognitive functioning, highlighting the importance of physical fitness ([Hillman et al., 2009a](#)). That is, the effect of exercise on the brain is modulated by the overall impact of physical exertion on the body. Therefore, not only is physical exercise necessary, but it should have specific characteristics that improve study participants' physical condition ([Hötting and Röder, 2013](#) ; [Reloba-Martínez et al., 2017](#) ; [Reigal et al., 2019a](#)).

Among the manifestations of physical fitness, aerobic capacity best explains the association between physical exercise and cognitive development in children and adolescents, as several authors have highlighted ([Pontifex et al., 2011](#) ; [Herting et al., 2014](#)).

Studies using neuroimaging techniques to explore these relationships have linked structural changes in the brain to exercise and the physical condition of children and adolescents. Authors such as [Chaddock et al. \(2010\)](#) have observed a higher volume in the hippocampus and the striated dorsal body with higher levels of aerobic fitness in children. Furthermore, [Chaddock-Heyman et al. \(2018\)](#) found an increase in the white matter microstructure of the genu of the corpus callosum after a 9-month program of moderate and vigorous physical activity in children. Likewise, in a group of obese children, [Esteban-Cornejo et al. \(2017\)](#) observed a relationship between cardiorespiratory capacity and speed/agility with the volume of gray matter in frontal, temporal, and subcortical regions, such as the premotor and supplementary motor cortex, the hippocampus, the caudate nucleus, as well as the inferior, superior, and parahippocampal temporal rotation.

Elderly People

The relationship between physical exercise and brain functioning has also been analyzed with great intensity in elderly people. There is a decline in certain physical and cognitive capacities that compromise normal functioning and autonomy during this stage of life ([Schiebener and Brand, 2017](#)). In addition, events such as retirement, the appearance of age-related illnesses, or the reduction in social relationships due to the loss of loved ones may occur, leading to greater isolation. In recent years, a number of studies have suggested that physical exercise in elderly people has benefits for aspects of brain functioning, such as attention, memory, or executive functioning ([Bherer et al., 2013](#) ; [Kramer and Colcombe, 2018](#) ; [Crespillo-Jurado et al., 2019](#)). Regular moderate to vigorous physical exercise has

been described as being protective against cognitive impairment and effective in maintaining adequate functioning in later life ([Zhu et al., 2017](#)). Therefore, promoting physical practice has become a potent strategy to improve elderly people's adaptation to the environment, maintaining, or improving their state of health and increasing their quality of life.

It has been observed that improvements in aspects such as aerobic capacity, balance, strength, or body composition would reduce the impact of aging on the deterioration of the brain, cushioning its effects and maintaining mental skills for longer periods of time ([Chang et al., 2012b](#) ; [Kerr et al., 2013](#) ; [Wilczyński et al., 2017](#)). As with other populations, cognitive functioning in elderly people is associated with their physical condition and changes in the structure of the brain support their functional development ([Reiter et al., 2015](#) ; [Fernandes et al., 2017](#)). For example, [Erickson et al. \(2011\)](#) observed that increases in aerobic capacity may be associated with improvements in spatial memory as well as increased BDNF concentration and volume in the hippocampus. [Boyle et al. \(2015\)](#) found that elderly people who were more active and had a lower body mass index had greater brain volume in areas such as the frontal and occipital lobes, in specific areas such as the frontal orbital cortex or the anterior cingulate gyrus, and had less dilation of cerebral ventricles.

In recent years, analyses of the impact of physical activity on neurodegenerative processes have received great attention. Several studies suggest that physical activity may be beneficial in response to conditions such as Alzheimer's disease, both delaying its onset, mitigating its

neuropathological effects, and improving brain functioning in patients ([Phillips et al., 2015](#)). Authors such as [Lautenschlager et al. \(2008\)](#) observed that, after a 24-week program of physical activity by elderly people, the rate of cognitive impairment in those at risk of dementia was reduced, suggesting that this type of intervention may inhibit the onset of the disease. It has also been observed that, in elderly people with a genetically increased risk of Alzheimer's disease, improvements in aspect so the physical conditions, such as cardiorespiratory capacity, may mitigate the risk of Alzheimer's disease ([Tari et al., 2019](#)). For this reason, it is considered that combined pharmacological treatments and more active lifestyles may be effective ways to counter this type of illness and other forms of dementia ([Mortimer and Stern, 2019](#)).

In addition, a large number of studies have indicated that physical activity may be useful in improving degenerative processes and symptoms in people with Parkinson's disease ([Fernández-del-Olmo et al., 2018](#)). In this population, it has been observed that the regular practice of physical activity throughout life could protect against the emergence of the disease, and that greater quantity and intensity have protective effects against Parkinson's disease ([Paillard et al., 2015](#)). Specifically, several authors have highlighted that aerobic exercise and cardiorespiratory capacity are positively related to brain processes, which again moderate the impact of Parkinson's disease ([Ahlskog, 2018](#)).

Limitations and Prospects for Future Research

Although many studies in recent years confirm the relationship between physical activity and cognitive functioning, there remains much to be <https://assignbuster.com/physical-activity-sports-practice-and-cognitive-functioning-the-current-research-status/>

understood. In addition, not all observed findings are supported by other studies, or they are based on small sample sizes with low statistical power. Thus, continued research is needed ([Frederiksen et al., 2018](#)). For greater precision in understanding the observed relationship between physical exercise and cognitive functioning and the possibilities for practical application, integration of knowledge from a range of scientific fields is recommended; these fields include sports science, neurobiology, neurophysiology, or neuropsychology as well as data from clinical and epidemiological studies. Furthermore, the transfer of findings from animal research to humans is required ([Tari et al., 2019](#)). According to recent research, there is some agreement on accepting the positive links observed between exercise and the brain, but the biological mechanisms underlying them require further examination ([Fernández-del-Olmo et al., 2018](#)).

Among the limitations of research to date is the heterogeneity of the studies and interventions, the lack of controls for strange variables, and small sample sizes. This has generated a collection of studies supporting the benefits of exercise on cognitive functioning, despite their moderate clinical value ([Brasure et al., 2018](#)). In addition, the exact amount and intensity of physical activity to meet individual needs has not yet been determined ([Paillard et al., 2015](#)). As an example, [Frederiksen et al. \(2018\)](#) , studying a group of elderly Alzheimer's patients, concluded that a 16-week (60 min/3 days per week) aerobic exercise program had no positive impact on cortical gray matter atrophy. However, disease symptoms improved and there were positive correlations in the intervention group between frontal cortical volume, exercise load, measures of attention, and cognitive processing

speed. Thus, the authors suggested that longer interventions and a larger sample could generate more convincing results. The challenge therefore remains to find the most appropriate formulas for each case and thereby determine the optimal exercise program.

An attempt has been made to explain the benefits of physical activity for the brain and its functioning in multiple populations, both healthy and with some pathology. However, presumably the requirements for each age or disease are specific, in addition to the differential impact on each person ([Phillips et al., 2015](#) ; [Carson et al., 2016](#) ; [Donnelly et al., 2016](#) ; [Fernandes et al., 2017](#)). For example, correlational or intervention studies in childhood and adolescence are conditioned by the natural development of these age groups and subject to various confounding factors, both intrapersonal and environmental, which impede the interpretation of the observed results ([Chen et al., 2016](#) ; [Chaddock-Heyman et al., 2018](#)). Thus, one of the greatest challenges in coming years is to identify the appropriate load, intensity, volume, frequency, duration, and type of physical activity so that the effects on the development of the brain of the relevant population group will achieve the desired objectives. Therefore, extensive control of confounding variables is necessary to minimize their bias. This goal achieved, not only can physical exercise for health be recommended in general terms, but it can also be implemented in multiple specific educational and clinical programs to complement the actions recommended by other disciplines.

Cognitive Functioning and High-Performance Sports

In the field of high-performance sports, the study of cognitive functioning has caught the attention of researchers. [Vestberg et al. \(2017\)](#) and [Policastro et al. \(2018\)](#) have shown that improved brain functioning in male and female athletes may enhance performance and predict success in competition. In general, it is assumed that this cognitive functioning may be more relevant in open sports requiring constant attention, management of multiple variables, or adaption to changing situations ([Williams et al., 2011](#) ; [Verburgh et al., 2014](#)). Furthermore, good cognitive functioning may be a competitive advantage in disciplines with less variability but requiring high levels of concentration or attentional control ([Memmert, 2009](#)).

Studies have observed that better scores in executive functioning are related to greater expertise and success in football players ([Verburgh et al., 2014](#) ; [Huijgen et al., 2015](#)). In turn, [Roca et al. \(2018\)](#) suggested that the creativity shown by adult football players in decision-making could be conditioned by their attentional capacity. In addition, [Voss et al. \(2010\)](#) pointed out that elite athletes tend to show better measures in aspects such as cognitive processing speed or various attentional tasks. Similarly, [Wagner et al. \(2014\)](#) concluded that cognitive aspects such as attentional capacity and executive functioning influence sports performance in handball. Moreover, [Hänggi et al. \(2015\)](#) pointed out that the brain structure can indicate the athlete's functioning, making brain plasticity an interesting predictor of player behavior on the playing field.

Recently, following these findings and the need to find variables to explain athletes' behaviors and success, the amount of research on how their brains work is increasing. Moreover, some researchers have attempted to train the athletes' brains with the objective of modifying the way athletes respond to stimuli during games and improve their decisions ([Calmels et al., 2004](#) ; [Seo et al., 2012](#)). Some of these studies have sought to modify the capacity of the athletes' brains and have tried to improve their attention, processing speed, or different aspects of cognitive functioning in sports such as tennis, football, handball, basketball, badminton, rugby, or archery ([Hagemann et al., 2006](#) ; [Ducrocq et al., 2016](#) ; [Romeas et al., 2016](#)).

For cognitive assessment and training in sports, multiple strategies have been used, both classical paper and pencil or computerized tests ([Memmert, 2009](#) ; [Hernández-Mendo et al., 2012](#) ; [Verburgh et al., 2014](#) ; [Huijgen et al., 2015](#) ; [Reigal et al., 2019b](#)) as well as other more technologically advanced procedures ([Memmert, 2009](#) ; [Hernández-Mendo et al., 2012](#) ; [Verburgh et al., 2014](#) ; [Huijgen et al., 2015](#)). In recent years, new electromechanical, digital, and combined devices have been developed, such as Fitlight Trainer™ , Dynavision D2™ , NeuroTracker™ , eye-tracking in sports, Vision Trainer™ , Senaptec Sensory Station, and Sports Vision Performance (M&S). Additionally, other products are based on technologies such as augmented or virtual reality. Overall, this new technology enabled notable progress in the assessment and cognitive preparation of athletes ([Schack et al., 2014](#) ; [Romeas et al., 2016](#) ; [Appelbaum and Erickson, 2018](#)). The rapid development of new technical measurement and training tools continuously increases the ecological validity of the measures by bringing the athlete's

<https://assignbuster.com/physical-activity-sports-practice-and-cognitive-functioning-the-current-research-status/>

training experience closer to an authentic game context. Therefore, the technology development opens up a very promising path for this line of research by offering possibilities that were previously unexplored.

Limitations and Prospects for Future Research

Studies of the relationship between brain functioning and sports performance have evolved rapidly in recent decades following technological advances that have enabled the development of increasingly powerful instruments ([Appelbaum and Erickson, 2018](#)). This has created a series of opportunities for professionals as well as researchers to advance our understanding of the relationship between brain functioning and sports performance. Similarly, current limitations are likely to be reduced by the coming technologies. Among them are many laboratory-based procedures used to study how the brain works in sport contexts ([Verburgh et al., 2014](#) ; [Fink et al., 2018](#)). This involves assessing the athletes' behavior in artificial situations, which limits the ecological validity of these procedures. Although possibly quite accurate inferences may be drawn, there are variables inherent in the competitive context that are currently difficult to reproduce.

Therefore, to increase the usefulness of cognitive assessments and training in performance sports, it will be necessary to improve the transference of laboratory knowledge to real-life contexts. The available technology allows experiences to approach reality, but such strategies are limited by technologies that are still very invasive and difficult to use on the playing field ([Romeas et al., 2016](#) ; [Appelbaum and Erickson, 2018](#)). As technical evolution makes it possible to evaluate and train brain function in sports

environments, it will help to determine more precisely the most appropriate way to stimulate the brain to optimize sports performance.

Similarly, cognitive assessments of athletes must be adjusted to the requirements of their specific tasks. In other words, the needs of a defender in football are not the same as those of a striker, nor are the needs of a basketball player the same as those of a tennis player. Therefore, another challenge to be addressed more clearly is finding the right type of training for each athlete considering his/her sporting characteristics and goals ([Vestberg et al., 2017](#)). In this way, the usefulness of this type of training, as well as the demand from athletes and technical coaches, will be increased. Much progress has been made in recent decades in areas such as physical or technical-tactical training. Nevertheless, a core question remains: how should the brain be trained to optimize sports performance so that it benefits physical and psychological health? Hopefully this question will be clearly answered in the future.

Author Contributions

AH-M, RR, JL-W, SS, OS, VM-S, RJ-R, JT-R, and CF participated in the design of the work and the bibliographic review, drafted the manuscript, and approved the final manuscript as submitted. All authors made substantial contributions to the final manuscript.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

<https://assignbuster.com/physical-activity-sports-practice-and-cognitive-functioning-the-current-research-status/>

References

- Ahlskog, J. E. (2018). Aerobic exercise: evidence for a direct brain effect to slow parkinson disease progression. *Mayo Clin. Proc.* 93, 360–372. doi: 10.1016/j.mayocp.2017.12.015
- Appelbaum, L. G., and Erickson, G. (2018). Sports vision training: a review of the state-of-the-art in digital training techniques. *Int. Rev. Sport Exerc. Psychol.* 11, 160–189. doi: 10.1080/1750984X.2016.1266376
- Becker, L., Kutz, D. F., and Voelcker-Rehage, C. (2016). Exercise induced changes in basal ganglia volume and their relation to cognitive performance. *J. Neurol. Neuromed.* 1, 19–24. doi: 10.29245/2572.942x/2016/5.1044
- Bherer, L., Erickson, K. I., and Liu-Ambrose, T. (2013). A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J. Aging Res.* 2013, 657508. doi: 10.1155/2013/657508
- Boecker, H., and Drzezga, A. (2016). A perspective on the future role of brain pet imaging in exercise science. *Neuroimage* 131, 73–80. doi: 10.1016/j.neuroimage.2015.10.021
- Boyle, C. P., Raji, C. A., Erickson, K. I., Lopez, O. L., Becker, J. T., Gach, H. M., et al. (2015). Physical activity, body mass index, and brain atrophy in Alzheimer's disease. *Neurobiol. Aging* 36, S194–S202. doi: 10.1016/j.neurobiolaging.2014.05.036
- Brasure, M., Desai, P., Davila, H., Nelson, V. A., Calvert, C., Jutkowitz, E., et al. (2018). Physical activity interventions in preventing cognitive decline and
- <https://assignbuster.com/physical-activity-sports-practice-and-cognitive-functioning-the-current-research-status/>

Alzheimer-type dementia: a systematic review. *Ann. Intern. Med.* 168, 30–38. doi: 10. 7326/M17-1528

Cabeza, R., Albert, M., Belleville, S., Craik, F. I., Duarte, A., Grady, C. L., et al. (2018). Maintenance, reserve and compensation: the cognitive neuroscience of healthy ageing. *Nat. Rev. Neurosci.* 19, 701–710. doi: 10. 1038/s41583-018-0068-2

Calmels, C., Berthoumieux, C., and d'Arripe-Longueville, F. (2004). Effects of an imagery training program on selective attention of national softball players. *Sport Psychol.* 18, 272–296. doi: 10. 1123/tsp. 18. 3. 272

Carson, V., Hunter, S., Kuzik, N., Wiebe, S. A., Spence, J. C., Friedman, A., et al. (2016). Systematic review of physical activity and cognitive development in early childhood. *J. Sci. Med. Sport* 19, 573–578. doi: 10. 1016/j. jsams. 2015. 07. 011

Chaddock, L., Erickson, K. I., Prakash, R. S., Kim, J. S., Voss, M. W., VanPatter, M., et al. (2010). A neuroimaging investigation of the association between aerobic fitness, hippocampal volume, and memory performance in preadolescent children. *Brain Res.* 1358, 172–183. doi: 10. 1016/j. brainres. 2010. 08. 049

Chaddock, L., Hillman, C. H., Buck, S. M., and Cohen, N. J. (2011). Aerobic fitness and executive control of relational memory in preadolescent children. *Med. Sci. Sports Exerc.* 43, 344–349. doi: 10. 1249/MSS. 0b013e3181e9af48

Chaddock-Heyman, L., Erickson, K. I., Kienzler, C., Drollette, E., Raine, L., Kao, S. C., et al. (2018). Physical activity increases white matter microstructure in children. *Front. Neurosci.* 12: 950. doi: 10. 3389/fnins. 2018. 00950

Chaddock-Heyman, L., Erickson, K. I., Voss, M. W., Knecht, A. M., Pontifex, M. B., Castelli, D. M., et al. (2013). The effects of physical activity on functional MRI activation associated with cognitive control in children: a randomized controlled intervention. *Front. Hum. Neurosci.* 7: 72. doi: 10. 3389/fnhum. 2013. 00072

Chang, Y. K., Labban, J. D., Gapin, J. I., and Etnier, J. L. (2012a). The effects of acute exercise on cognitive performance: a meta-analysis. *Brain Res.* 1453, 87–101. doi: 10. 1016/j. brainres. 2012. 02. 068

Chang, Y. K., Pan, C. Y., Chen, F. T., Tsai, C. L., and Huang, C. C. (2012b). Effect of resistance-exercise training on cognitive function in healthy older adults: a review. *J. Aging Phys. Act.* 20, 497–517. doi: 10. 1123/japa. 20. 4. 497

Chen, A. G., Zhu, L. N., Yan, J., and Yin, H. C. (2016). Neural basis of working memory enhancement after acute aerobic exercise: fMRI study of preadolescent children. *Front. Psychol.* 7: 1804. doi: 10. 3389/fpsyg. 2016. 01804

Cheron, G., Petit, G., Cheron, J., Leroy, A., Cebolla, A., Cevallos, C., et al. (2016). Brain oscillations in sport: toward EEG biomarkers of performance. *Front. Psychol.* 7: 246. doi: 10. 3389/fpsyg. 2016. 00246

Costigan, S. A., Eather, N., Plotnikoff, R. C., Hillman, C. H., and Lubans, D. R. (2016). High-intensity interval training for cognitive and mental health in adolescents. *Med. Sci. Sport Exerc.* 48, 1985–1993. doi: 10.1249/MSS.0000000000000993

Crespillo-Jurado, M., Delgado-Giralt, J., Reigal, R. E., Rosado, A., Wallace-Ruiz, A., Juárez Ruiz de Mier, R., et al. (2019). Body composition and cognitive functioning in a sample of active elders. *Front. Psychol.* 10: 1569. doi: 10.3389/fpsyg.2019.01569

Dalen, T., Sandmael, S., Stevens, T. G., Hjelde, G. H., Kjøsnes, T. N., and Wisløff, U. (2019). Differences in acceleration and high-intensity activities between small-sided games and peak periods of official matches in elite soccer players. *J. Strength Cond. Res.* doi: 10.1519/JSC.0000000000003081 [Epub ahead of print].

Donnelly, J. E., Tomporowski, P. D., Hillman, C. H., Castelli, D. M., Etnier, J. L., Lee, S. M., et al. (2016). physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Med. Sci. Sport Exerc.* 48, 1197–1222. doi: 10.1249/MSS.0000000000000901

Ducrocq, E., Wilson, M., Vine, S., and Derakshan, N. (2016). Training attentional control improves cognitive and motor task performance. *J. Sport Exerc. Psychol.* 38, 521–533. doi: 10.1123/jsep.2016-0052

Dumoulin, S. O., Fracasso, A., van der Zwaag, W., Siero, J. C., and Petridou, N. (2018). Ultra-high field MRI: advancing systems neuroscience towards

mesoscopic human brain function. *Neuroimage* 168, 345–357. doi: 10.1016/j.neuroimage.2017.01.028

Elleberg, D., and St-Louis-Deschênes, M. (2010). The effect of acute physical exercise on cognitive function during development. *Psychol. Sport Exerc.* 11, 122–126. doi: 10.1016/j.psychsport.2009.09.006

Erickson, K. I., Voss, M. W., Prakash, R. S., Basak, C., Szabo, A., Chaddock, L., et al. (2011). Exercise training increases size of hippocampus and improves memory. *Proc. Natl. Acad. Sci. U. S. A.* 108, 3017–3022. doi: 10.1073/pnas.1015950108

Erickson, K. I., Hillman, C. H., and Kramer, A. F. (2015). Physical activity, brain, and cognition. *Curr. Opin. Behav. Sci.* 4, 27–32. doi: 10.1016/j.cobeha.2015.01.005

Esteban-Cornejo, I., Cadenas-Sanchez, C., Contreras-Rodriguez, O., Verdejo-Roman, J., Mora-Gonzalez, J., Migueles, J. H., et al. (2017). A whole brain volumetric approach in overweight/obese children: examining the association with different physical fitness components and academic performance. The Active Brains project. *Neuroimage* 159, 346–354. doi: 10.1016/j.neuroimage.2017.08.011

Fernandes, J., Arida, R. M., and Gómez-Pinilla, F. (2017). Physical exercise as an epigenetic modulator of brain plasticity and cognition. *Neurosci. Biobehav. R.* 80, 443–456. doi: 10.1016/j.neubiorev.2017.06.012

Fernández-del-Olmo, M., Sánchez-Molina, J. A., Morenilla, L., Gómez-Varela, J., Fernández-Lago, H., Bello, O., et al. (2018). Aerobic and resistance exercises in Parkinson's disease: a narrative review. *Eur. J. Hum. Mov.* 41, 149–174.

Fink, A., Rominger, C., Benedek, M., Perchtold, C. M., Papousek, I., Weiss, E. M., et al. (2018). EEG alpha activity during imagining creative moves in soccer decision-making situations. *Neuropsychologia* 114, 118–124. doi: 10.1016/j.neuropsychologia.2018.04.025

Fister, I., Fister, D., Deb, S., Mlakar, U., and Brest, J. (2018). Post hoc analysis of sport performance with differential evolution. *Neural Comput. Appl.* 1–10. doi: 10.1007/s00521-018-3395-3

Fontes, E. B., Okano, A. H., De Guio, F., Schabort, E. J., Min, L. L., Basset, F. A., et al. (2015). Brain activity and perceived exertion during cycling exercise: an fMRI study. *Br. J. Sport Med.* 49, 556–560. doi: 10.1136/bjsports-2012-091924

Frederiksen, K. S., Larsen, C. T., Hasselbalch, S. G., Christensen, A. N., Høgh, P., Wermuth, L., et al. (2018). A 16-week aerobic exercise intervention does not affect hippocampal volume and cortical thickness in mild to moderate Alzheimer's disease. *Front. Aging Neurosci.* 10: 293. doi: 10.3389/fnagi.2018.00293

Gale, C. R., Cooper, R., Craig, L., Elliott, J., Kuh, D., Richards, M., et al. (2012). Cognitive function in childhood and lifetime cognitive change in relation to

mental wellbeing in four cohorts of older people. *PLoS One* 7: e44860. doi: 10.1371/journal.pone.0044860

Gutmann, B., Zimmer, P., Hülsdünker, T., Lefebvre, J., Binnebößel, S., Oberste, M., et al. (2018). The effects of exercise intensity and post-exercise recovery time on cortical activation as revealed by EEG alpha peak frequency. *Neurosci. Lett.* 668, 159–163. doi: 10.1016/j.neulet.2018.01.007

Hagemann, N., Strauss, B., and Cañal-Bruland, R. (2006). Training perceptual skill by orienting visual attention. *J. Sport Exerc. Psychol.* 28, 143–158. doi: 10.1123/jsep.28.2.143

Hänggi, J., Langer, N., Lutz, K., Birrer, K., Mérillat, S., and Jäncke, L. (2015). Structural brain correlates associated with professional handball playing. *PLoS One* 10: e0124222. doi: 10.1371/journal.pone.0124222

Henriksen, K., Storm, L. K., Stambulova, N., Pyrdol, N., and Larsen, C. H. (2019). Successful and less successful interventions with youth and senior athletes: insights from expert sport psychology practitioners. *J. Clin. Sport Psychol.* 13, 72–94. doi: 10.1123/jcsp.2017-0005

Hernández-Mendo, A., Martínez Jiménez, M. A., Pastrana Brincones, J. L., and Morales Sánchez, V. (2012). Computer program for evaluation and training of care. *Rev. Iberoam. Psicol. Ejercicio Deport.* 7, 339–358.

Herting, M. M., Colby, J. B., Sowell, E. R., and Nagel, B. J. (2014). White matter connectivity and aerobic fitness in male adolescents. *Dev. Cogn. Neurosci.* 7, 65–75. doi: 10.1016/j.dcn.2013.11.003

Hillman, C. H., Buck, S. M., Themanson, J. R., Pontifex, M. B., and Castelli, D. M. (2009a). Aerobic fitness and cognitive development: event-related brain potential and task performance indices of executive control in preadolescent children. *Dev. Psychol.* 45, 114–129. doi: 10.1037/a0014437

Hillman, C. H., Pontifex, M. B., Raine, L. B., Castelli, D. M., Hall, E. E., and Kramer, A. F. (2009b). The effect of acute treadmill walking on cognitive control and academic achievement in preadolescent children. *Neuroscience* 159, 1044–1054. doi: 10.1016/j.neuroscience.2009.01.057

Hillman, C. H., Castelli, D. M., and Buck, S. M. (2005). Aerobic fitness and neurocognitive function in healthy preadolescent children. *Med. Sci. Sport Exerc.* 37, 1967–1974. doi: 10.1249/01.mss.0000176680.79702.ce

Hillman, C. H., Pontifex, M. B., Castelli, D. M., Khan, N. A., Raine, L. B., Scudder, M. R., et al. (2014). Effects of the FITKids randomized controlled trial on executive control and brain function. *Pediatrics* 134, e1063–e1071. doi: 10.1542/peds.2013-3219

Hötting, K., and Röder, B. (2013). Beneficial effects of physical exercise on neuroplasticity and cognition. *Neurosci. Biobehav. R.* 37, 2243–2257. doi: 10.1016/j.neubiorev.2013.04.005

Hsu, C. L., Best, J. R., Davis, J. C., Nagamatsu, L. S., Wang, S., Boyd, L. A., et al. (2018). Aerobic exercise promotes executive functions and impacts functional neural activity among older adults with vascular cognitive impairment. *Br. J. Sport Med.* 52, 184–191. doi: 10.1136/bjsports-2016-096846

Huang, P., Fang, R., Li, B. Y., and Chen, S. D. (2016). Exercise-related changes of networks in aging and mild cognitive impairment brain. *Front. Aging Neurosci.* 8: 47. doi: 10.3389/fnagi.2016.00047

Huijgen, B. C., Leemhuis, S., Kok, N. M., Verburgh, L., Oosterlaan, J., Elferink-Gemser, M. T., et al. (2015). Cognitive functions in elite and sub-elite youth soccer players aged 13 to 17 years. *PLoS One* 10: e0144580. doi: 10.1371/journal.pone.0144580

Hynynen, S. T., Van Stralen, M. M., Sniehotta, F. F., Araújo-Soares, V., Hardeman, W., Chinapaw, M. J., et al. (2016). A systematic review of school-based interventions targeting physical activity and sedentary behaviour among older adolescents. *Int Rev Sport Exerc. Psychol.* 9, 22–44. doi: 10.1080/1750984X.2015.1081706

Jonasson, L. S., Nyberg, L., Kramer, A. F., Lundquist, A., Riklund, K., and Boraxbekk, C. J. (2017). Aerobic exercise intervention, cognitive performance, and brain structure: results from the physical influences on brain in aging (PHIBRA) study. *Front. Aging Neurosci.* 8: 336. doi: 10.3389/fnagi.2016.00336

Kerr, J., Marshall, S. J., Patterson, R. E., Marinac, C. R., Natarajan, L., Rosenberg, D., et al. (2013). Objectively measured physical activity is related to cognitive function in older adults. *J. Am. Geriatr. Soc.* 61, 1927–1931. doi: 10.1111/jgs.12524

Kramer, A. F., and Colcombe, S. (2018). Fitness effects on the cognitive function of older adults: a meta-analytic study—revisited. *Perspect. Psychol. Sci.* 13, 213–217. doi: 10.1111/1467-9280.t01-1-01430

Krenn, B., Finkenzeller, T., Würth, S., and Amesberger, G. (2018). Sport type determines differences in executive functions in elite athletes. *Psychol. Sport Exerc.* 38, 72–79. doi: 10.1016/j.psychsport.2018.06.002

Lautenschlager, N. T., Cox, K. L., Flicker, L., Foster, J. K., van Bockxmeer, F. M., Xiao, J., et al. (2008). Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *JAMA* 300, 1027–1037. doi: 10.1001/jama.300.9.1027

Li, J. W., O'Connor, H., O'Dwyer, N., and Orr, R. (2017). The effect of acute and chronic exercise on cognitive function and academic performance in adolescents: a systematic review. *J. Sci. Med. Sport* 20, 841–848. doi: 10.1016/j.jsams.2016.11.025

Lourenco, M. V., Frozza, R. L., Zhang, H., Kincheski, G. C., Ribeiro, F. C., Gonçalves, R. A., et al. (2019). Exercise-linked FNDC5/irisin rescues synaptic plasticity and memory defects in Alzheimer's models. *Nat. Med.* 25, 165–175. doi: 10.1038/s41591-018-0275-4

Lubans, D., Richards, J., Hillman, C., Faulkner, G., Beauchamp, M., Nilsson, M., et al. (2016). Physical activity for cognitive and mental health in youth: a systematic review of mechanisms. *Pediatrics* 138: e20161642. doi: 10.1542/peds. 2016-1642

Memmert, D. (2009). Pay attention! A review of visual attentional expertise in sport. *Int. Rev. Sport Exerc. Psychol.* 2, 119–138. doi: 10.1080/17509840802641372

Moran, A., Campbell, M., and Toner, J. (2019). Exploring the cognitive mechanisms of expertise in sport: progress and prospects. *Psychol. Sport Exerc.* 42, 8–15. doi: 10.1016/j.psychsport. 2018. 12. 019

Mortimer, J. A., and Stern, Y. (2019). Physical exercise and activity may be important in reducing dementia risk at any age. *Neurology* 92: 8. doi: 10.1212/WNL. 0000000000006935

Northey, J. M., Cherbuin, N., Pumpa, K. L., Smee, D. J., and Rattray, B. (2018). Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br. J. Sport Med.* 52, 154–160. doi: 10.1136/bjsports-2016-096587

Paillard, T., Rolland, Y., and de Souto Barreto, P. (2015). Protective effects of physical exercise in Alzheimer's disease and Parkinson's disease: a narrative review. *J. Clin. Neurol.* 11, 212–219. doi: 10.3988/jcn. 2015. 11. 3. 212

- Pérez-Lobato, R., Reigal, R. E., and Hernández-Mendo, A. (2016). Relationships between physical practice, physical condition, and attention in a sample of adolescents. *J. Sport Psychol.* 25, 179–186.
- Phillips, C., Baktir, M. A., Das, D., Lin, B., and Salehi, A. (2015). The link between physical activity and cognitive dysfunction in Alzheimer disease. *Phys. Ther.* 95, 1046–1060. doi: 10. 2522/ptj. 20140212
- Policastro, F., Accardo, A., Marcovich, R., Pelamatti, G., and Zoia, S. (2018). Relation between motor and cognitive skills in italian basketball players aged between 7 and 10 Years Old. *Sports* 6: E80. doi: 10. 3390/sports6030080
- Pontifex, M. B., McGowan, A. L., Chandler, M. C., Gwizdala, K. L., Parks, A. C., Fenn, K., et al. (2019). A primer on investigating the after effects of acute bouts of physical activity on cognition. *Psychol. Sport Exerc.* 40, 1–22. doi: 10. 1016/j. psychsport. 2018. 08. 015
- Pontifex, M. B., Raine, L. B., Johnson, C. R., Chaddock, L., Voss, M. W., Cohen, N. J., et al. (2011). Cardiorespiratory fitness and the flexible modulation of cognitive control in preadolescent children. *J. Cogn. Neurosci.* 23, 1332–1345. doi: 10. 1162/jocn. 2010. 2152
- Reigal, R. E., Barrero, S., Martín, I., Morales-Sánchez, V., Juárez-Ruiz de Mier, R., and Hernández-Mendo, A. (2019a). Relationships between reaction time, selective attention, physical activity, and physical fitness in children. *Front. Psychol.* 10: 2278. doi: 10. 3389/fpsyg. 2019. 02278

Reigal, R. E., González-Guirval, F., Morillo-Baro, J. P., Morales-Sánchez, V., Juárez-Ruiz de Mier, R., and Hernández-Mendo, A. (2019b). Effects of a computerized training on attentional capacity of young soccer players. *Front. Psychol.* 10: 2279. doi: 10.3389/fpsyg.2019.02279

Reiter, K., Nielson, K. A., Smith, T. J., Weiss, L. R., Alfini, A. J., and Smith, J. C. (2015). Improved cardiorespiratory fitness is associated with increased cortical thickness in mild cognitive impairment. *J. Int. Neuropsychol. Soc.* 21, 757–767. doi: 10.1017/S135561771500079X

Reloba-Martínez, S., Reigal, R. E., Hernández-Mendo, A., Martínez-López, E. J., Martín-Tamayo, I., and Chirrosa-Ríos, L. J. (2017). Effects of vigorous extracurricular physical exercise on the attention of schoolchildren. *J. Sports Psychol.* 26, 29–36.

Roca, A., Ford, P. R., and Memmert, D. (2018). Creative decision making and visual search behavior in skilled soccer players. *PLoS One* 13: e0199381. doi: 10.1371/journal.pone.0199381

Romeas, T., Guldner, A., and Faubert, J. (2016). 3D-Multiple Object Tracking training task improves passing decision-making accuracy in soccer players. *Psychol. Sport Exerc.* 22, 1–9. doi: 10.1016/j.psychsport.2015.06.002

Sakamoto, S., Takeuchi, H., Ihara, N., Ligao, B., and Suzukawa, K. (2018). Possible requirement of executive functions for high performance in soccer. *PLoS One* 13: e0201871. doi: 10.1371/journal.pone.0201871

Schack, T., Bertollo, M., Koester, D., Maycock, J., and Essig, K. (2014). "Technological advancements in sport psychology," in *Routledge Companion to Sport and Exercise Psychology*, eds A. G. Papaioannou, and D. Hackfort, (London: Routledge), 953–965.

Schiebener, J., and Brand, M. (2017). Age-related variance in decisions under ambiguity is explained by changes in reasoning, executive functions, and decision-making under risk. *Cogn. Emot.* 31, 816–824. doi: 10.1080/02699931.2016.1159944

Schwarb, H., Johnson, C. L., Daugherty, A. M., Hillman, C. H., Kramer, A. F., Cohen, N. J., et al. (2017). Aerobic fitness, hippocampal viscoelasticity, and relational memory performance. *Neuroimage* 153, 179–188. doi: 10.1016/j.neuroimage.2017.03.061

Scudder, M. R., Federmeier, K. D., Raine, L. B., Direito, A., and Boyd, J. K. (2014). The association between aerobic fitness and language processing in children: implications for academic achievement. *Brain Cogn.* 87, 140–152. doi: 10.1016/j.bandc.2014.03.016

Seo, J., Kim, Y. T., Song, H. J., Lee, H. J., Lee, J., Jung, T. D., et al. (2012). Stronger activation and deactivation in archery experts for differential cognitive strategy in visuospatial working memory processing. *Behav. Brain Res.* 229, 185–193. doi: 10.1016/j.bbr.2012.01.019

Shih, C. H., Moore, K., Browner, N., Sklerov, M., and Dayan, E. (2019). Physical activity mediates the association between striatal dopamine

transporter availability and cognition in Parkinson's disease. *Parkinsonism Relat. Disord.* 62, 68–72. doi: 10.1016/j.parkreldis.2019.01.027

Sorrells, S. F., Paredes, M. F., Cebrian-Silla, A., Sandoval, K., Qi, D., Kelley, K. W., et al. (2018). Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555, 377–381. doi: 10.1038/nature25975

Tari, A. R., Norevik, C. S., Scrimgeour, N. R., Kibro-Flatmoen, A., Storm-Mathisen, J., Bergersen, L. H., et al. (2019). Are the neuroprotective effects of exercise training systemically mediated? *Prog. Cardiovasc. Dis.* 62, 94–101. doi: 10.1016/j.pcad.2019.02.003

Verburgh, L., Scherder, E. J., van Lange, P. A., and Oosterlaan, J. (2014). Executive functioning in highly talented soccer players. *PLoS One* 9: e91254. doi: 10.1371/journal.pone.0091254

Vestberg, T., Reinebo, G., Maurex, L., Ingvar, M., and Petrovic, P. (2017). Core executive functions are associated with success in young elite soccer players. *PLoS One* 12: e0170845. doi: 10.1371/journal.pone.0170845

Voss, M. W., Kramer, A. F., Basak, C., Prakash, R. S., and Roberts, B. (2010). Are expert athletes 'expert' in the cognitive laboratory? A meta-analytic review of cognition and sport expertise. *Appl. Cogn. Psychol.* 24, 812–826. doi: 10.1002/acp.1588

- Voss, M. W., Soto, C., Yoo, S., Sodoma, M., Vivar, C., and van Praag, H. (2019). Exercise and hippocampal memory systems. *Trends Cogn. Sci.* 23, 318–333. doi: 10. 1016/j. tics. 2019. 01. 006
- Wagner, H., Finkenzeller, T., Würth, S., and von Duvillard, S. P. (2014). Individual and team performance in team-handball: a review. *J. Sport Sci. Med.* 13, 808–816.
- Wenner, C. J., Bianchi, J., Figueredo, A. J., Rushton, J., and Jacobs, W. J. (2013). Life history theory and social deviance: the mediating role of executive function. *Intelligence* 41, 102–113. doi: 10. 1016/j. intell. 2012. 11. 004
- Wilczyński, J., Pedrycz, A., Mucha, D., Ambroży, T., and Mucha, D. (2017). Body posture, postural stability, and metabolic age in patients with Parkinson's disease. *Bio. Med. Res. Int.* 2017: 3975417. doi: 10. 1155/2017/3975417
- Williams, A. M., Ford, P., Eccles, D. W., and Ward, P. (2011). Perceptual-cognitive expertise in sport and its acquisition: implications for applied cognitive psychology. *Appl. Cogn. Psychol.* 25, 432–442. doi: 10. 1002/acp. 1710
- Xue, Y., Yang, Y., and Huang, T. (2019). Effects of chronic exercise interventions on executive function among children and adolescents: a systematic review with meta-analysis. *Br. J. Sport Med.* 53, 1397. doi: 10. 1136/bjsports-2018-099825

Zhu, W., Wadley, V. G., Howard, V. J., Hutto, B., Blair, S. N., and Hooker, S. P. (2017). Objectively Measured Physical Activity and Cognitive Function in Older Adults. *Med. Sci. Sport Exerc.* 49, 47–53. doi: 10.1249/MSS.0000000000001079

Zylke, J., and Bauchner, H. (2016). The unrelenting challenge of obesity. *JAMA* 315, 2277–2278. doi: 10.1001/jama.2016.6190