

Occlusal stent construction research



**ASSIGN
BUSTER**

Initial examination was carried out consisting of evaluation the periodontal condition of the teeth. After selecting the suitable patients for the study, all of them received supra gingival scaling and polishing with a good motivation and instruction in oral hygiene measures including brushing, using dental floss and interproximal brushes as indicated and demonstration was given to them about the work of perio Q gel and it's application .

An alginate impression was taken and an occlusal stent was constructed for each patient. After completion of the occlusal stent construction, the patient was recalled again and this was considered the first visit. In this visit the clinical periodontal examination was carried out for the selected sites and it included the following parameters:

- Plaque Index (PLI) :- (Silness and Loe 1964)

A periodontal probe was used after air drying of the teeth and the selected sites were examined for plaque. The periodontal probe was gently passed along the gingival crevice. The criteria was the following:

Score 0: No plaque in the gingival area.

Score 1: A film of plaque adhering to the free gingival margin and adjacent area of the tooth surface, which cannot be seen with the naked eye but only by using disclosing solution or by using probe.

Score 2: Moderate accumulation of soft deposits within the gingival pocket, on the gingival margin and or adjacent tooth surface which can be seen by naked eye.

Score 3: Abundance of soft matter within the gingival pocket and or on the gingival margin and adjacent tooth.

- Gingival Index(GI): (*Loe, 1967*).

The occurrence of gingival inflammation was assessed using the criteria of gingival index system

Score 0: Normal gingiva.

Score 1: Mild inflammation, slight change in color, slight edema, and no bleeding on probing

Score 2: Moderate inflammation, redness and glazing, bleeding on probing.

Score 3: Severe inflammation, marked redness and ulceration, tendency to spontaneous bleeding.

- Bleeding on Probing (BOP) :- (*Carranza, 2012*).

A blunt periodontal probe inserted to the bottom of the periodontal pocket/sulcus and is moved gently along the root surface. If bleeding occurred within 30 seconds after probing, the site was given positive score (1), and a negative score (0) for the non-bleeding site

- Probing Pocket Depth (PPD):(*Lindhe et al., 1998*)

The probing pocket depth was measured with a William's periodontal probe at four sites of all teeth on (mesial, buccal, distal and lingual), the distance from gingival margin to the most apical extent of the probe inserted parallel to the long axis of the tooth to the nearest millimeter (mm) was recorded only for the sites exhibiting probing depth of (5-8)mm.

- Relative Attachment Level (RAL):

The occlusal stent was adjusted to fit the teeth, then vertical grooves or holes corresponding to the probed site were made using rotary fissure bur, these grooves provided a fixed reference mark for probe insertion and angulation. The stent was putted on the occlusal surfaces to cover half or 2/3 of the crown. The distance

from the base of the pocket to the lower border of the stent at the base of the groove was considered as the RAL. The measurement was made to the nearest mm .

The clinical periodontal recordings were repeated after 3 and 6 weeks.

Treatments

After recording of all periodontal parameters for the selected sites, the patient mouth was splitted into three quadrants, each quadrant received different treatment modality and as follows:

Initial visit (1st day): patient selection, supra gingival scaling, alginate impression, motivation, instruction.

. Gel group: 111 sites in this group received intra pocket application of perio Q gel. The selected sites were isolated by cotton rolls and dried the teeth by air, and then dried the pockets by paper point size (30, 35, 40, and 45). the application of the gel was made using disposable syringe of 5ml. the sharp tip of needle was removed by rotary bur to avoid hurting the gingival tissue and smoothed it, then 1 ml of the gel was pulled by the syringe and the needle gently placed down through the pocket until it reach to the bottom of

the pockets then placed the gel while worked the way up until the gingival margin. Each pocket was received a range of (0. 1-0. 3) ml., the excess gel oozing from the pockets was removed by Cotton rolls The patients were instructed to avoid spitting, washing, eating and drinking for 2 hours of the gel application. Toothbrush and interdental aids should paused of the day after the gel application.

Clinical

periodont

al

First paramete

visit rs

(1day) recorded,

gel

applicatio

n.

Secon Clinical

d visit periodont

(after al

3 paramete

weeks rs

) recorded

Third Clinical

visit periodont

(after al

paramete
6week
rs
s)
recorded

Combination group: 106 sites in this group received scaling and root planing, then after one hour, the patient examined if there was no blood oozing, then the gel applied as was described previously. If not, the patient was referred to the next day.

Clinical
periodont
al
paramete
rs
recorded,

First root
visit planning
(1day) to the
selected
sites,
after 1
hour
putted
the gel.

Secon Clinical
d visit periodont

(after al
3 paramete
weeks rs
) recorded

Clinical
Third periodont
visit al
(after paramete
6week rs
s) recorded

Scaling and root planning group: 106 sites in this group received scaling and root planning only.

Clinical
periodont
al
paramete
First rs
visit recorded,
(1day) root
planning
to the
selected
sites.

Second Clinical

3rd visit periodont

(after al

3 parameters

weeks

) recorded

Third Clinical

3rd visit periodont

(after al

6 weeks parameters

6 weeks

rs

s)

recorded

Pilot study

To perform intra examiner calibration and inter examiner calibration with clinical periodontal parameters used in this study (PLI, GI, BOP, PPD, RALI), a pilot study was carried out in Department of periodontics, College of Dentistry, Baghdad University. It was carried out about four weeks before the commencement of the actual project on two subjects with twelve sites. The intra examiner calibration was repeated after an appropriate period (usually 2-4 weeks) to resolve any memory bias. While the inter examiner calibration was repeated by another trained professional at the same time. The consistency (calibration) should be at least 90% and if it is low, the measurement should be repeated.

Statistical Analysis

<https://assignbuster.com/occlusal-stent-construction-research/>

Data were processed and analyzed using SPSS 16 for windows8 (statistical package for social science) and excel 2013. both descriptive and inferential analyses

Descriptive Statistics

- Tables (Range, Frequencies and Percentage)
- Arithmetic Mean
- Standard Deviation
- Mean Difference.
- Median
- Minimum and Maximum
- Graphical Presentation by Bar Charts and Scattered Plots.

Inferential Statistics

There was used to accept or reject the statistical hypotheses, which included:

- Analysis of Variance Test(ANOVA) One Way
- Student t-Test for equality of means of two independent groups.
- Wilcoxon Signs rank test
- Mann Whitney U Test

References

- Lindhe J Karring T Lang N. Clinical periodontology and implant dentistry. 3rd edition. Copenhagen, Munksgaard, 1998

- Loe, H. The Gingival Index, the Plaque Index and the Retention Index Systems. *Journal of Periodontology*, Vol. 38, No. 6 (November-December 1967), pp. 610-616.
- Silness J, Loe H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand* 1964; 22: 112-135