

# Environmental enrichment and successful aging

[Health & Medicine](#)



## Introduction

In more than a dozen animal models of human neurological pathology, environmental enrichment has been shown to ameliorate the human-like neurological symptoms present in these animals models, including the cognitive decline in aging ( [Segovia et al., 2006](#) ; [Patel, 2012](#) ) and in Alzheimer's disease ( [Arendash et al., 2004](#) ; [Jankowsky et al., 2005](#) ; [Lazarov et al., 2005](#) ; [Berardi et al., 2007](#) ; [Costa et al., 2007](#) ; [Basak et al., 2008](#) ; [Arranz et al., 2011](#) ; [Polito et al., 2014](#) ).

[Humes et al. \(2013\)](#) suggested that the multi-sensory decline that accompanies normal aging in humans might influence the decline in cognition. They systematically examined changes in hearing, vision, touch, and cognitive function, in young, middle, and older adults. By varying the difficulty and sensitivity of the tasks, as well as making allowances for deteriorating sensory skills, they not only were able to show that age, global sensory, and global cognitive abilities were correlated, but also they further suggested that the decline in sensory processing and acuity may affect the cognitive decline. We present here a brief summary of the deterioration of the sensory and cognitive systems that occurs in older adults and we will present evidence that suggests that enriching the environment of the aging brain by reversing the decline in sensory systems may aid cognitive functioning.

## Hearing

### Neural Deterioration With Hearing Loss

Age-related hearing loss is common in older adults, with almost two-thirds of those over 70 years old having compromised hearing, and about a third of older adults experiencing debilitating hearing loss ([Lin et al., 2011b](#); [WHO, 2018](#)). The odds of hearing loss are 5. 5-fold higher in men than in women ([Agrawal et al., 2008](#)) and hearing sensitivity declines two times faster in men than in women, with diminished sensitivity observed in males as early as age 30 years ([Syka, 2002](#)).

Changes in brain anatomy accompany the decrements in auditory functioning. A reduction in the volume of gray matter is observed bilaterally in auditory cortex that correlates with the loss of high-frequency age-related hearing loss ([Peelle et al., 2011](#); [Eckert et al., 2012](#)). There also are neural losses in areas that do not process auditory information directly. [Husain et al. \(2011\)](#) found that older adults with hearing loss experienced decreases in the volume of gray matter in the superior and medial frontal gyri, both of which are involved in cognitive processes, along with losses in the superior temporal cortex, which includes the auditory cortex and multisensory association cortex, and right anterior cingulate cortex, which includes both emotional and cognitive areas. [Wong et al. \(2010\)](#) found that older adults who had limitations in their ability to perceive speech in a noisy environment had smaller frontal cortices than younger adults with normal auditory ability.

In a longitudinal study over 6 years, [Lin et al. \(2014\)](#) compared whole-brain volumes of older adults with hearing impairment to those without hearing loss and found that those with hearing loss had an accelerated loss of brain

<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

tissue. There were also volume losses in the right temporal lobe, which included the superior, middle, and inferior temporal gyri, and the parahippocampus, areas that also mediate cognitive processes.

As for functional changes in the brain in response to auditory signals in hearing-impaired individuals, differences in evoked responses correlated with hearing impairment in noisy environments ([Campbell and Sharma, 2013](#)). In noisy environments, older adults with hearing loss also respond to speech with both a reduction in the activity of auditory brain areas and an increase in the activity of parietofrontal brain centers ([Wong et al., 2010](#)). In general, additional cognitive resources are recruited to achieve auditory perception under challenging auditory conditions for older adults ([Eckert et al., 2008](#); [Harris et al., 2009](#); [Erb and Obleser, 2013](#); [Erb et al., 2013](#); [Vaden et al., 2013, 2016](#)).

### Cognitive Deterioration With Hearing Loss

There is a consistent finding that has been revealed in five recent meta-analyses that hearing impairment in older adults increases the risk of cognitive deficiencies, including dementia ([Thomson et al., 2017](#); [Wei et al., 2017](#); [Zheng et al., 2017](#); [Loughrey et al., 2018](#); [Yuan et al., 2018](#)).

Indeed, the risk of cognitive loss increased by 29% compared to normal-hearing older adults over the course of follow-ups of up to 6 years and that risk was increased by 57% when the patients were followed for a longer period, even when potential confounding factors were considered ([Yuan et al., 2018](#)). Moreover, increased peripheral damage of the auditory system predicted a greater risk of cognitive impairment in older adults ([Yuan et al., 2018](#)). Older adults with mild to severe hearing impairment have a two- to

five-fold increase in their risk of developing dementia compared to older adults with normal hearing ([Lin et al., 2011a](#)). These cognitive impairments lead to a decreased ability to care for themselves, as confirmed by the impaired performance on the activities of daily living scales in older adults with hearing loss ([Gopinath et al., 2013](#)).

There are a number of ways that hearing impairment could affect cognition. The first is that there is a common mechanism that affects both neural systems independently, such as a gradual circulatory impairment, although the predictive value of hearing impairment appears to occur years before the cognitive decline ([Albers et al., 2015](#)). Another possibility is that the loss of hearing makes it difficult to have social interactions and increases the risk of depression, both of which could negatively impact cognitive ability ([Mick et al., 2014](#); [Dawes et al., 2015](#)). Also possible is that as more effort is required for the listening process, fewer resources may be available for comprehension and memory, whereas the easing of the listening effort results in improvements in cognitive performance ([Pichora-Fuller, 2006](#)). Finally, it may be that the loss of auditory stimulation deprives the cognitive areas of their normal level of stimulation and thereby allows those areas to deteriorate.

### **Enhanced Auditory Stimulation and Cognitive Function**

If the loss of auditory stimulation allows the brain's cognitive systems to deteriorate, would either the use of hearing aids or cochlear implants reverse the cognitive loss? Indeed, many studies have shown that the use of hearing aids is associated with improved cognitive abilities and decreased depression in older adults who have hearing loss ([Mulrow et al., 1990](#); [Tesch-Römer,](#)

[1997](#) ; [Acar et al., 2011](#) ; [Choi et al., 2011](#) ; [Dawes et al., 2015](#) ; [Castiglione et al., 2016](#) ; although see [van Hooren et al., 2005](#) ).

Hearing aids, however, simply amplify and filter sounds and do not recreate the lost frequencies for their wearers. On the other hand, cochlear implants, even in older adults, restore a significant portion of the sound range that had been lost to the individual ( [Williamson et al., 2009](#) ; [Carlson et al., 2010](#) ). Such implants restore not only auditory function, but also restore social interactions and quality of life for hearing impaired older adults ( [Olze et al., 2012](#) ). In addition, the cochlear implants induced improvements in cognitive function. Specifically, 12 months after cochlear implantation plus auditory training, 81% of the subjects with the lowest cognitive scores had improved their cognitive function ( [Mosnier et al., 2015](#) ). In another study of cochlear implantation in older adults with hearing impairment, the cochlear implants resulted in improvements in 70% of all cognitive tests, although the subjects did worse on 25% of the cognitive tests after surgery ( [Cosetti et al., 2016](#) ). Social isolation and depression also improved in those with auditory rehabilitation ( [Castiglione et al., 2016](#) ).

Interestingly, enhanced auditory enrichment started early in life also appears to impart benefits to cognition in later life. Lifelong music training appears to ameliorate the cognitive decline observed in older adults ( [Hanna-Pladdy and MacKay, 2011](#) ; [Parbery-Clark et al., 2012](#) ; [Zendel and Alain, 2012](#) ).

Moreover, older adults with musical experience have enhanced ability to process speech ( [Bidelman and Alain, 2015](#) ). [White-Schwoch et al. \(2013\)](#) further showed that even moderate amounts of early music training had a

persistent effect on maintaining the cognitive ability of older adults, even with a lapse of decades since they had played their instrument.

## Vision

### Visual and Cognitive Deterioration in Older Adults

Vision is compromised with age ([Klaver et al., 1998](#); [Muñoz et al., 2000](#); [Salvi et al., 2006](#)) and there is an increased risk of cognitive loss, Alzheimer's disease and an increased clinical severity of Alzheimer's disease among those with visual impairment ([Uhlmann et al., 1991](#); [Lin et al., 2004](#); [Reyes-Ortiz et al., 2005](#); [Whitson et al., 2010, 2012](#); [Ong et al., 2012](#); [Chen et al., 2013, 2017](#); [Mine et al., 2016](#)). Indeed, older adults with good visual acuity have a 63% decreased risk for developing dementia over an 8.5-year period and older adults with diminished visual acuity are 5 times more likely to experience cognitive loss than are older adults with good vision ([Rogers and Langa, 2010](#)). In addition, older adults with age-related cataracts have a 1.4 times increased likelihood of developing Alzheimer's disease ([Lai et al., 2014](#)). It is also the case that older adults with poor vision engage in fewer activities that involve their cognitive ability than those with normal vision ([Varin et al., 2017](#)).

It is possible that there is a common mechanism that causes both vision and cognition to deteriorate. In fact, [Drobny et al. \(2005\)](#) found that older adults with poor vision did poorly on tasks that did not involve vision, suggesting that general cognition is deteriorating as vision deteriorates. When [Dickinson and Rabbitt \(1991\)](#) simulated visual impairment in normal subjects, the subjects were able to read the passage accurately, but their recall of material suffered, suggesting that the difficulty in storing the memory

interfered with its retrieval. The cognitive loss that accompanies visual loss therefore may be due to such a situation.

### **Neural Deterioration With Compromised Vision**

Older adults with macular degeneration causing their visual impairment have a smaller visual cortex, along with a thinner cortex, smaller surface area, and lower gray matter volume than older adults with normal vision ( [Hernowo et al., 2014](#) ; [Prins et al., 2016](#) ). [Chen et al. \(2013\)](#) found deterioration of gray matter in the visual cortical areas in those individuals with open-angle glaucoma, along with finding increased gray matter in some cognitive areas, including the medial area of the superior parietal cortex. The size of the frontal lobe and other areas involved in cognition decreases in older adults with macular degeneration, along with deterioration of the connections between visual cortex and frontal cortex, suggesting a cognitive link in this situation ( [Chen et al., 2013](#) ; [Hernowo et al., 2014](#) ).

Blind individuals also have a significantly larger anterior right hippocampus compared to that of normal sighted individuals, and the posterior right hippocampus is significantly smaller in blind individuals ( [Leporé et al., 2009](#) ), suggesting modifications to regions of the brain involved with cognitive processing in response to diminished visual information.

### **Compromised/Rectified Visual Stimulation, Cognitive Ability, and Visual Cortex**

Wearing reading glasses is associated with improved cognitive outcomes for older adults ( [Spierer et al., 2016](#) ), although this condition could simply be the result of an increased ability to read. Indeed, the difference between groups was diminished when education level was considered. Unilateral

cataract surgery in older adults can result in improved visual acuity, and that surgery improves cognitive functioning ([Tamura et al., 2004](#); [Ishii et al., 2008](#); [Miyata et al., 2016](#)), or maintains cognitive functioning (in comparison with other groups, which showed worsened cognitive functioning across the duration of the study; [Elliott et al., 2009](#)). It should be noted, however, that [Anstey et al. \(2006\)](#) did not find cognitive benefits after cataract surgery. [Jefferis et al. \(2015\)](#) found a small improvement in cognition after cataract surgery that was not correlated with improvement in visual acuity.

Older adults with visual impairments have a decrease in the volume of their visual cortex ([Boucard et al., 2009](#)) and unilateral cataract surgery can result in both an improvement in vision and an increase in visual cortex gray matter contralateral to the operated eye ([Lou et al., 2013](#)). [Lin et al. \(2018\)](#) found both improvements in visual acuity, cognitive functioning, and brain activity following bilateral cataract surgery, and these improvements were accompanied by an increase in gray matter volume in visual cortex, 6-months post-surgery.

### **Combined Hearing and Visual Impairment**

Diminished functionality in both vision and hearing senses, or dual sensory impairment (DSI), is a common problem in the older adult population, with greater than 70% of individuals with major vision impairments also exhibiting hearing loss ([Heine and Browning, 2002](#)). Individuals with both visual and hearing impairment have an even higher risk of cognitive loss, a diminished capacity on the daily living scales measures, an increased incidence of depression, a greater likelihood of social isolation, an increased risk of falls,

and an increased risk of mortality ( [Keller et al., 1999](#) ; [Saunders and Echt, 2007](#) ; [Gopinath et al., 2013, 2016](#) ; [Mitoku et al., 2016](#) ). Indeed, when combined, visual and auditory acuity account for 49. 2% of the total variance and 93. 1% of the age-related variance in intelligence in older adults ( [Lindenberger and Baltes, 1994](#) ). Both the high prevalence and the detrimental effects of DSI in the older adult population have resulted in recommendations for treatment that include mechanical aids to improve both hearing and vision deficits ( [Saunders and Echt, 2007](#) ; [Zhou and Faure Walker, 2015](#) ; [Gopinath et al., 2016](#) ).

## Olfaction

### Neural Deterioration With Olfactory Loss

Normal human aging is accompanied by a deterioration of olfactory abilities ( [Hoffman et al., 2016](#) ; [Dong et al., 2017](#) ; [Seubert et al., 2017](#) ), with 18% of older adults having a significant olfactory impairment and 46% of those over 80 years old having very limited olfactory ability ( [Hoffman et al., 1998](#) ; [Murphy et al., 2002](#) ; [Doty et al., 2015](#) ; [Toussaint et al., 2015](#) ; [Liu et al., 2016](#) ).

Differing from other sensory systems, which have cortical projections that are gated by the thalamus, the olfactory system has direct projections to brain regions involved with cognition and the loss or compromise of the olfaction system results in volume losses to many of these cortical brain areas in humans at any age ( [Bitter et al., 2010a, b, 2011](#) ; [Peng et al., 2013](#) ; [Shen et al., 2013](#) ; [Yao et al., 2014](#) ). Olfactory projection sites, which include primary and secondary sensory cortical areas, as well as cortical

regions involved with cognitive processing, also deteriorate with age ( [Segura et al., 2013](#) ; [Kollendorfer et al., 2015](#) ).

### **Olfactory System Deterioration and Cognitive Loss**

Olfactory dysfunction also accompanies or precedes the early symptoms of cognitive disorders such as Alzheimer's disease, Parkinson's disease, Lewy body dementia, frontotemporal dementia, Creutzfeldt-Jakob disease, mild cognitive impairment, and schizophrenia ( [Doty et al., 1988](#) ; [Devanand et al., 2000](#) ; [Ponsen et al., 2004](#) ; [Tabaton et al., 2004](#) ; [Ross et al., 2006](#) ; [Luzzi et al., 2007](#) ; [Wattendorf et al., 2009](#) ; [Devanand et al., 2010](#) ; [Li et al., 2010](#) ; [Meusel et al., 2010](#) ; [Nguyen et al., 2010](#) ; [Parrao et al., 2012](#) ; [Sohrabi et al., 2012](#) ; [Conti et al., 2013](#) ). Given that these cognitive disorders have widely differing etiologies, it raises the possibility that the loss of olfactory stimulation contributes to the decline in cognitive ability in some of these disorders, particularly those in which olfactory loss precedes cognitive loss. Moreover, a degradation of olfactory ability predicts an elevated risk of mild cognitive impairment (MCI) and the degree of olfactory degradation may be used to predict which individuals with MCI will develop Alzheimer's disease ( [Devanand et al., 2000](#) ; [Peters et al., 2003](#) ; [Schubert et al., 2008, 2017](#) ; [Roberts et al., 2016](#) ; [Lafaille-Magnan et al., 2017](#) ; [Adams et al., 2018](#) ).

### **Enhanced Olfactory Stimulation**

Increasing olfactory stimulation in individuals who have experienced olfactory loss due to a variety of problems, such as post-infectious olfactory dysfunction, head trauma, Parkinson's disease, and aging, has been shown to improve olfactory identification, olfactory discrimination, and to a lesser extent, olfactory thresholds ( [Hummel et al., 2009](#) ; [Haehner et al., 2013](#) ;

[Konstantinidis et al., 2013](#) ; [Damm et al., 2014](#) ; [Geißler et al., 2014](#) ; [Patel et al., 2017](#)). These results were achieved using twice daily fragrance exposures to four odorants taken from each of four odor groups: resinous (eucalyptus), flowery (rose), fruity (lemon), and aromatic (clove) and the individuals continued this regimen for varying durations, (i. e., 12 weeks to 6 months). Further improvements in olfactory ability were observed with increased duration of exposure, increased concentration of the odorants, or an increased number of odorants ( [Damm et al., 2014](#) ; [Altundag et al., 2015](#) ; [Konstantinidis et al., 2016](#) ). A recent review and meta-analysis by [Pekala et al. \(2016\)](#) concluded that olfactory training was effective in improving olfactory system function (e. g., odor discrimination and identification), and positive results were observed following the loss of olfactory function with different etiologies. In addition to improvements in sensory ability, older adults exposed to increased olfactory stimulation have an improvement in their cognitive function, as evidenced by increased verbal fluency, an improvement in their depressive symptoms, and an improved sense of wellbeing ( [Birte-Antina et al., 2018](#) ).

## Mastication

### Cognitive Deterioration With Dental Problems

There are hundreds of reports that have examined the relationship between mastication (chewing) and cognition. Researchers have found a strong correlation between cognitive decline and poor oral health (for reviews, see [Chen et al., 2015](#) ; [Azuma et al., 2017](#) ). For example, there are a number of studies showing that the fewer teeth an individual has, the worse their cognitive ability is ( [Bergdahl et al., 2007](#) ; [Stein et al., 2007](#) ; [Syrjälä et al., 2007](#) ).

[2007](#); [Grabe et al., 2009](#); [Okamoto et al., 2010](#); [Lexomboon et al., 2012](#); [Del Brutto et al., 2014](#); [Mummolo et al., 2014](#); [Peres et al., 2014](#); [Elsig et al., 2015](#); [Luo et al., 2015](#); [Stewart et al., 2015](#)). Contributing factors that can lead to tooth loss are periodontal disease ([Noble et al., 2009](#); [Kamer et al., 2012](#); [Gil-Montoya et al., 2015](#); [Welmer et al., 2017](#)), untreated tooth caries ([Tonetti et al., 2017](#)), and low socioeconomic scale ([Cabrera et al., 2005](#); [Matsuyama et al., 2017](#)). The subsequent consequences of tooth loss include increased stress levels ([Budtz-Jørgensen, 1980](#)), decreased social interactions, lower quality of life ([Griffin et al., 2012](#)), and a limited diet ([Walls et al., 2000](#); [Spaccavento et al., 2009](#); [Kimura et al., 2013](#)). All of these factors may be involved in the degradation of cognition, making it difficult to conclude that chewing supports cognition directly.

On the other hand, when rats and mice have teeth removed, or are given only powdered or liquid food to eat, their cognitive ability also declines ([Kato et al., 1997](#); [Yamamoto and Hirayama, 2001](#); [Fukushima-Nakayama et al., 2017](#)), along with a decrease in brain-derived neurotrophic factor (BDNF), synaptic density, and neuronal number in the hippocampus ([Onozuka et al., 1999](#); [Yamamoto and Hirayama, 2001](#); [Okihara et al., 2014](#); [Takeda et al., 2016](#)). In addition, there is widespread volumetric loss of gray matter in the brain following tooth loss in mice ([Avivi-Arber et al., 2016](#)). At the same time, disruption of normal chewing leads to elevated levels of corticosterone ([Kubo et al., 2007](#)) and lower levels of hippocampal glucocorticoid receptors ([Ichihashi et al., 2008](#)) that are associated with chronic stress ([Sapolsky et al., 1984](#)). The advantage of examining chewing activity in non-human species is that the caloric intake and amount of food can be equalized across

the hard and soft food eating conditions, thus maintaining similar nutritional values. The results of these animal studies support the idea that the active process of chewing plays a role in cognitive functioning and stress reduction. Indeed, in humans, a stronger correlation was observed between chewing ability (with or without the use of dental prostheses) and cognitive function than that observed between tooth loss and cognition ([Lexomboon et al., 2012](#)).

### **Enhanced Mastication/Corrected Dentition and Cognitive Restoration**

While the loss of normal chewing ability is associated with cognitive decline, increased chewing activity has been shown to decrease stress and improve cognition in humans ([Baker et al., 2004](#); [Stephens and Tunney, 2004](#); [Scholey et al., 2009](#); [Ono et al., 2010](#); [Yu et al., 2013](#); [Azuma et al., 2017](#); for a review, see [Weijenberg et al., 2011](#)). Chewing gum alone can increase attention, decrease reaction times, and improve mood, even under stressful conditions ([Smith, 2010](#); [Kubo et al., 2015](#)), as well as increasing cognitive processing speed ([Hirano et al., 2013](#)). A meta-analysis found positive effects of chewing on alertness or attention in 64% of the studies they examined ([Hirano and Onozuka, 2015](#)). In contrast, [Tucha et al. \(2004\)](#) observed chewing gum only changed aspects of attention without cognitive improvement. Consistent with the notion that chewing may aid cognition, the brain areas that are activated by chewing include cognitive centers such as the frontal cortex and the medial temporal lobe ([Onozuka et al., 2002](#); [Choi et al., 2017](#)).

Rats and mice given wood dowels to chew have a reduced stress response and are able to maintain their hippocampal-dependent cognitive function (<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

[Ono et al., 2010](#) ; [Miyake et al., 2012](#) ; [Chen et al., 2015](#)). In addition, the chewing action relieves the stress-induced suppression of cell proliferation in the hippocampal dentate gyrus that may underlie the impaired hippocampal functioning with tooth loss ([Kubo et al., 2007](#)). Also, switching mice from powdered food to hard pellet food reverses the suppression of neurogenesis in the forebrain subventricular zone that occurred while on the soft food diet ([Utsugi et al., 2014](#)).

Humans who have lost teeth and were subsequently given dentures or dental implants that restored dental function also had improved prefrontal cortex activity ([Narita et al., 2009](#) ; [Kimoto et al., 2011](#) ; [Kamiya et al., 2016](#)), along with improved cognitive performance ([Cerutti-Kopplin et al., 2015](#) ; [Banu et al., 2016](#) ; [De Cicco et al., 2016](#)).

## Sensorimotor/Somatosensory Stimulation

### Exercise Increases/Restores the Size of Cortical Structures

Decreases in the size of the brain, including the hippocampus, are observed in normally aging older adults, and this decrease accompanies the declines in cognitive functioning ([Raz et al., 2005](#)). Aerobic exercise can restore some of these decreases in brain volume. About 12 months of aerobic exercise resulted in an increase in the volume of the hippocampus ([Erickson et al., 2011, 2014](#) ; [Niemann et al., 2014](#)) and 6 months of aerobic training increased the volume of gray and white matter, while stretching and toning for the same amount of time did not result in brain volume changes ([Colcombe et al., 2006](#)). In addition, both hippocampal volume and dorsolateral prefrontal cortex thickness were positively correlated with aerobic fitness ([Jonasson et al., 2017](#)).

## Cognitive Benefits of Exercise

Multiple studies, reviews, and meta-analyses have concluded that in older adults, exercise training results in improvements in cognitive skills, along with improved health and mobility ([Colcombe and Kramer, 2003](#); [Hertzog et al., 2008](#); [Kemoun et al., 2010](#); [Bherer et al., 2013](#); [Kirk-Sanchez and McGough, 2014](#); [Barha et al., 2017](#); [Gregory et al., 2017](#); [Kennedy et al., 2017](#); [Mavros et al., 2017](#); [Saez de Asteasu et al., 2017](#); [Northeby et al., 2018](#); for review see [Erickson et al., 2013](#)). In addition, following exercise training, improvements in cognitive functioning are correlated with measurable increases in brain volume ([Erickson et al., 2011](#); [Mortimer et al., 2012](#)) and increased BDNF levels ([Erickson et al., 2011](#); [Sungkarat et al., 2018](#)), both of which may underlie the observed cognitive gains ([Erickson et al., 2013](#)). Exercise training also restores the decrease in interhemispheric inhibition typically observed in older adults, which could lead to improved motor control ([McGregor et al., 2018](#)). In addition, using fMRI to measure brain activity, exercise for 12 months improved functional connectivity among the cortices and the improved connectivity was associated with improved executive function ([Voss et al., 2010](#)). Moreover, adults who exercised showed altered activity in the right inferior frontal gyrus during a semantic verbal fluency task that was correlated with improvements in the task ([Nocera et al., 2017](#)). A recent meta-analysis that examined the influence of exercise, cognitive training, or the two combined on falls and cognition in older adults with MCI found that exercise or the combination of exercise and cognitive training resulted in gait speed and balance improvements, as well as cognitive function gains, all of which can contribute to decreased incidence of falls ([Lipardo et al., 2017](#)).

## Mechanisms by Which Exercise Can Improve Cognition

[Stimpson et al. \(2018\)](#) have proposed a model for the mechanism underlying cognitive improvement following exercise. Specifically, they first note that exercise increases cerebral angiogenesis and circulation in the brain. [Pereira et al. \(2007\)](#) reported that after 3 months of aerobic exercise, there was increased neurogenesis and increased cerebral blood volume in the human hippocampus, raising the possibility that increased angiogenesis mediated these improvements. There is also a transient elevation of serum BDNF after humans exercise ([Ferris et al., 2007](#); [Håkansson et al., 2017](#)), and when the action of BDNF is blocked in rats after exercise, the cognitive benefits are also blocked ([Vaynman et al., 2004](#)). They next noted that exercise also decreases chronic inflammation in older adults ([Cotman et al., 2007](#)) and that restores elevated levels of insulin-like growth factor (IGF-1), which elevates BDNF levels and increases neurogenesis ([Carro et al., 2000](#)). The increased neurogenesis then allows the improved cognitive outcomes following exercise. It should be noted, however, that the presence of adult human neurogenesis has been recently questioned ([Sorrells et al., 2018](#)).

It is also the case that a relatively recent meta-analysis on studies showing aerobic gains with exercise came to the conclusion that there was no compelling evidence indicating that exercise improves cognitive abilities in older adults ([Young et al., 2015](#)). If increased aerobic capacity does not drive the effects of exercise, then it seems possible that the increase in somatosensory stimulation that is experienced during exercise may drive any improvements that are seen following exercise. This perspective would

then bring any cognitive improvements with exercise in line with the effects of other types of sensory stimulation that we have discussed.

## Conclusion

### Deterioration of Sensory Systems Can Be Mitigated

It seems clear that the deterioration of sensorimotor systems contributes to the decline in cognition seen in older adults, either directly, by providing less stimulation to the cognitive areas of the brain, or indirectly, by depriving older adults of the nutrition, intellectual engagement, or social engagement that they need to thrive. Currently, there are methods that attempt to restore the declining systems, and although they are not always able to completely restore sensory system function, they do appear to aid individuals, with improvements to sensory acuity, cognition, and quality of life, thus improving chances of successful aging. Importantly, it is apparent that early detection of deficits to any of these systems is critical, and restoration or repair of any deficits should be prioritized to increase the likelihood of maintaining cognitive and full body health. Most certainly, new technologies are needed to continue to refine and improve the restoration of individual sensorimotor systems. Additional environmental enrichment using improvements to mastication or exercise training results in cognitive and health gains, which may potentially further aid in the successful aging of older adults.

### Barriers to Maintenance or Restoration of Cognition Function by Restoring Deteriorating Sensory Systems in Older Adults

Although the aids described previously for use in the restoration of sensory and motor systems are promising, the benefits are not without a cost and

may not be experienced by everyone. While it makes sense to do what one can to sustain or repair these systems, the cost of such repairs is not affordable by most people. Perhaps the greatest need is in hearing aids because a set can cost thousands of dollars and these devices are typically not covered by medical insurance in the United States. Consequently, only 3–4% of those with mild hearing loss who need hearing aids are wearing them ([Chien and Lin, 2012](#)). Cochlear implants, while even more expensive, can be covered by medical insurance, but are called for only in those with severe hearing impairments and in deaf individuals ([Lin et al., 2012](#)). The use of reading glasses seems to be a useful tool to allow older adults to engage with their environment effectively. While some investigators find a cognitive benefit for cataract surgery, others do not observe such an improvement, suggesting that the cognitive benefits may not be reliable. Dentures and dental implants are effective for maintaining or restoring cognitive ability for older adults with poor dentition, but again it comes at great expense, and it is something that Medicare does not cover. Enhanced olfactory stimulation has had limited tests for cognitive improvement and there are no studies showing improved neural responses after enhanced olfactory stimulation in older adults.

It is difficult to induce older adults to exercise regularly. Only about 16% of older adults engage in the recommended amount of physical activity on a daily basis ([Centers for Disease Control and Prevention \[CDC\], 2018](#)), which are at least 150 min of moderate-intensity aerobic activity or 75 min of vigorous-intensity aerobic activity and 2 or more days of muscle-strengthening activities per week ([Centers for Disease Control and](#)

[Prevention \[CDC\], 2018](#)). Indeed, 33% of older adults reported no physical activity at all ([Centers for Disease Control and Prevention \[CDC\], 2018](#)). Barriers to exercise include lack of motivation, lack of knowledge, pain, poor health, physical limitations, peer pressure, and bad weather ([Costello et al., 2011](#); [Gothe and Kendall, 2016](#)). Surprisingly, older adults do not have an increased risk of injury from exercise ([Stathokostas et al., 2013](#)). Although there is a concern that the oxidative stress associated with intense exercise would hasten aging in older adults, physically active older adults actually have reduced exercise-induced oxidative stress than older adults with a lower level of physical activity. Additionally, regular physical activity apparently improves the antioxidant defenses of older adults ([Meijer et al., 2002](#)).

### **The Possibility of a Common Factor**

While each sensory system may have its impact on cognition in its own way, there is also the possibility that there is a common factor that underlies their role in the maintenance of cognitive processes. For example, the loss or degradation of each sensory system may result in an emotional change that could impact cognitive performance. As discussed above, depression is associated with the loss of teeth ([Shah et al., 2015](#)), is associated with the loss of olfactory stimulation ([Kohli et al., 2016](#)), and often follows the degradation of the auditory and visual systems. Depression also often follows cognitive dysfunction in older adults ([Yin et al., 2015](#)). Alternatively, the common factor could be a physiological element that deteriorates both sensory and cognitive brain systems concurrently, but deficits are more easily detectable in the sensory systems. It also may be that continual

exposure to environmental elements may wear on the sensory systems before they impact the cognitive brain areas.

## Future Directions

One future direction may be to enhance stimulation of olfactory, visual, auditory, tactile, masticatory, and motor systems simultaneously or in conjunction. This type of sensorimotor stimulation had benefits for individuals with developmental neurological disorders ([Woo and Leon, 2013](#); [Woo et al., 2015](#); [Aronoff et al., 2016](#)). Environmental enrichment has also been shown to improve the symptoms of children with Rett syndrome ([Downs et al., 2018](#)). Similar forms of sensorimotor combination therapy may have health benefits for aging adults.

## Author Contributions

ML and CW conceived and wrote this review.

## Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## References

Acar, B., Yurekli, M. F., Babademez, M. A., Karabulut, H., and Karasen, R. M. (2011). Effects of hearing aids on cognitive functions and depressive signs in elderly people. *Arch. Gerontol. Geriatr.* 52, 250–252. doi: 10.1016/j.archger.2010.04.013

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Adams, D. R., Kern, D. W., Wroblewski, K. E., McClintock, M. K., Dale, W., and Pinto, J. M. (2018). Olfactory dysfunction predicts subsequent dementia in older U. S. adults. *J. Am. Geriatr. Soc.* 66, 140–144. doi: 10.1111/jgs.15048

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Agrawal, Y., Platz, E. A., and Niparko, J. K. (2008). Prevalence of hearing loss and differences by demographic characteristics among US adults: data from the National Health and Nutrition Examination Survey, 1999–2004. *Arch. Intern. Med.* 168, 1522–1530. doi: 10.1001/archinte.168.14.1522

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Albers, M. W., Gilmore, G. C., Kaye, J., Murphy, C., Wingfield, A., Bennett, D. A., et al. (2015). At the interface of sensory and motor dysfunctions and Alzheimer's disease. *Alzheimers Dement.* 11, 70–98. doi: 10.1016/j.jalz.2014.04.514

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Altundag, A., Cayonu, M., Kayabasoglu, G., Salihoglu, M., Tekeli, H., Saglam, O., et al. (2015). Modified olfactory training in patients with postinfectious olfactory loss. *Laryngoscope* 125, 1763–1766. doi: 10.1002/lary.25245

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Anstey, K. J., Lord, S. R., Hennessy, M., Mitchell, P., Mill, K., and von Sanden, C. (2006). The effect of cataract surgery on neuropsychological test performance: a randomized controlled trial. *J. Int. Neuropsychol. Soc.* 12, 632–639. doi: 10.1017/S1355617706060954

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Arendash, G. W., Garcia, M. F., Costa, D. A., Cracchiolo, J. R., Wefes, I. M., and Potter, H. (2004). Environmental enrichment improves cognition in aged Alzheimer's transgenic mice despite stable beta-amyloid deposition. *Neuroreport* 15, 1751–1754. doi: 10.1097/01.wnr.0000137183.68847.4e

[CrossRef Full Text](#) | [Google Scholar](#)

Aronoff, E., Hillyer, R., and Leon, M. (2016). Environmental enrichment therapy for autism: outcomes with increased access. *Neural Plast.* 2016: 2734915. doi: 10.1155/2016/2734915

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Arranz, L., De Castro, N. M., Baeza, I., Giménez-Llort, L., and De la Fuente, M. (2011). Effect of environmental enrichment on the immunoendocrine aging of male and female triple-transgenic 3xTg-AD mice for Alzheimer's disease. *J. Alzheimers Dis.* 25, 727–737. doi: 10.3233/JAD-2011-110236

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Avivi-Arber, L., Seltzer, Z., Friedel, M., Lerch, J. P., Moayedi, M., Davis, K. D., et al. (2016). Widespread volumetric brain changes following tooth loss in female mice. *Front. Neuroanat.* 10: 121. doi: 10.3389/fnana.2016.00121

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Azuma, K., Zhou, Q., Niwa, M., and Kubo, K. Y. (2017). Association between mastication, the hippocampus, and the HPA Axis: a comprehensive review. *Int. J. Mol. Sci.* 18: E1687. doi: 10.3390/ijms18081687

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Baker, J. R., Bezance, J. B., Zellaby, E., and Aggleton, J. P. (2004). Chewing gum can produce context-dependent effects upon memory. *Appetite* 43, 207–210. doi: 10.1016/j.appet.2004.06.004

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Banu, R. F., Veeravalli, P. T., and Kumar, V. A. (2016). Comparative evaluation of changes in brain activity and cognitive function of edentulous patients, with dentures and two-implant supported mandibular overdenture-pilot study. *Clin. Implant Dent. Relat. Res.* 18, 580–587. doi: 10.1111/cid.12336

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Barha, C. K., Davis, J. C., Falck, R. S., Nagamatsu, L. S., and Liu-Ambrose, T. (2017). Sex differences in exercise efficacy to improve cognition: a systematic review and meta-analysis of randomized controlled trials in older humans. *Front. Neuroendocrinol.* 46: 71–85. doi: 10.1016/j.yfrne.2017.04.002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Basak, C., Boot, W. R., Voss, M. W., and Kramer, A. F. (2008). Can training in a real-time strategy videogame attenuate cognitive decline in older adults? *Psychol. Aging* 23, 765–777. doi: 10.1111/cid.12336

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Berardi, N., Braschi, C., Capsoni, S., Cattaneo, A., and Maffei, L. (2007). Environmental enrichment delays the onset of memory deficits and reduces neuropathological hallmarks in a mouse model of Alzheimer-like neurodegeneration. *J. Alzheimers Dis.* 11, 359–370. doi: 10.3233/JAD-2007-11312

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bergdahl, M., Habib, R., Bergdahl, J., Nyberg, L., and Nilsson, L. G. (2007). Natural teeth and cognitive function in humans. *Scand. J. Psychol.* 48, 557–565. doi: 10.1111/j.1467-9450.2007.00610.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bherer, L., Erickson, K. I., and Liu-Ambrose, T. (2013). A review of the effects of physical activity and exercise on cognitive and brain functions in older adults. *J. Aging Res.* 2013: 657508. doi: 10.1155/2013/657508

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bidelman, G. M., and Alain, C. (2015). Musical training orchestrates coordinated neuroplasticity in auditory brainstem and cortex to counteract age-related declines in categorical vowel perception. *J. Neurosci.* 35, 1240–1249. doi: 10.1523/JNEUROSCI.3292-14.2015

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Birte-Antina, W., Ilona, C., Antje, H., and Thomas, H. (2018). Olfactory training with older people. *Int. J. Geriatr. Psychiatry* 33, 212–220. doi: 10.1002/gps. 4725

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bitter, T., Brüderle, J., Gudziol, H., Burmeister, H. P., Gaser, C., and Guntinas-Lichius, O. (2010a). Gray and white matter reduction in hyposmic subjects - A voxel-based morphometry study. *Brain Res.* 1347, 42–47. doi: 10.1016/j.brainres.2010.06.003

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bitter, T., Gudziol, H., Burmeister, H. P., Mentzel, H.-J., Guntinas-Lichius, O., and Gaser, C. (2010b). Anosmia leads to a loss of gray matter in cortical brain areas. *Chem. Senses* 35, 407–415. doi: 10.1093/chemse/bjq028

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Bitter, T., Siegert, F., Gudziol, H., Burmeister, H. P., Mentzel, H. J., Hummel, T., et al. (2011). Gray matter alterations in parosmia. *Neuroscience* 177, 177–182. doi: 10.1016/j.neuroscience.2011.01.016

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Boucard, C. C., Hernowo, A. T., Maguire, R. P., Jansonius, N. M., Roerdink, J. B., Hooymans, J. M., et al. (2009). Changes in cortical grey matter density

associated with long-standing retinal visual field defects. *Brain* 132, 1898–1906. doi: 10.1093/brain/awp119

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Budtz-Jørgensen, E. (1980). A 3-month study in monkeys of occlusal dysfunction and stress. *Scand. J. Dent. Res.* 88, 171–180. doi: 10.1111/j.1600-0722.1980.tb01211.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Cabrera, C., Hakeberg, M., Ahlqvist, M., Wedel, H., Björkelund, C., Bengtsson, C., et al. (2005). Can the relation between tooth loss and chronic disease be explained by socioeconomic status? A 24-year follow-up from the population study of women in Gothenburg, Sweden. *Eur. J. Epidemiol.* 20, 229–236. doi: 10.1007/s10654-004-5961-5

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Campbell, J., and Sharma, A. (2013). Compensatory changes in cortical resource allocation in adults with hearing loss. *Front. Syst. Neurosci.* 7: 71. doi: 10.3389/fnsys.2013.00071

[CrossRef Full Text](#) | [Google Scholar](#)

Carlson, M. L., Breen, J. T., Gifford, R. H., Driscoll, C. L., Neff, B. A., Beatty, C. W., et al. (2010). Cochlear implantation in the octogenarian and nonagenarian. *Otol. Neurotol.* 31, 1343–1349. doi: 10.1097/MAO.0b013e3181edb69d

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Carro, E., Nuñez, A., Busiguina, S., and Torres-Aleman, I. (2000). Circulating insulin-like growth factor I mediates effects of exercise on the brain. *J. Neurosci.* 20, 2926–2933. doi: 10.1523/JNEUROSCI.20-08-02926. 2000

[CrossRef Full Text](#) | [Google Scholar](#)

Castiglione, A., Benatti, A., Velardita, C., Favaro, D., Padoan, E., Severi, D., et al. (2016). Aging, cognitive decline and hearing loss: effects of auditory rehabilitation and training with hearing aids and cochlear implants on cognitive function and depression among older adults. *Audiol. Neurotol.* 21(Suppl. 1), 21–28. doi: 10.1159/000448350

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Centers for Disease Control and Prevention [CDC] (2018). *Nutrition, Physical Activity, and Obesity Data, Trends, and Maps*. Available at: [https://nccd.cdc.gov/NPAO\\_DTM/](https://nccd.cdc.gov/NPAO_DTM/)

[Google Scholar](#)

Cerutti-Kopplin, C., Emami, E., Hilgert, J. B., Hugo, F. N., and Padilha, D. M. P. (2015). Cognitive status of edentate elders wearing complete dentures: does quality of denture matter? *J. Dent.* 43, 1071–1075. doi: 10.1016/j.jdent.2015.07.008

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Chen, H., Iinuma, M., Onozuka, M., and Kubo, K. Y. (2015). Chewing maintains hippocampus-dependent cognitive function. *Int. J. Med. Sci.* 12, 502–509. doi: 10.7150/ijms.11911

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Chen, S. P., Bhattacharya, J., and Pershing, S. (2017). Association of vision loss with cognition in older adults. *JAMA Ophthalmol.* 135, 963–970. doi: 10.1001/jamaophthalmol.2017.2838

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Chen, W. W., Wang, N., Cai, S., Fang, Z., Yu, M., Wu, Q., et al. (2013). Structural brain abnormalities in patients with primary open-angle glaucoma: a study with 3T MR imaging. *Invest. Ophthalmol. Vis. Sci.* 54, 545–554. doi: 10.1167/iovs.12-9893

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Chien, W., and Lin, F. R. (2012). Prevalence of hearing aid use among older adults in the United States. *Arch. Intern. Med.* 172, 292–293. doi: 10.1001/archinternmed.2011.1408

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Choi, A. Y., Shim, H. J., Lee, S. H., Yoon, S. W., and Joo, E. J. (2011). Is cognitive function in adults with hearing impairment improved by the use of hearing aids? *Clin. Exp. Otorhinolaryngol.* 4, 72–76. doi: 10.3342/ceo.2011.4.2.72

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Choi, Y. H., Jang, W. H., Im, S. U., Song, K. B., Lee, H. K., Lee, H. D., et al. (2017). The brain activation pattern of the medial temporal lobe during chewing gum: a functional MRI study. *Neural Regen. Res.* 12, 812–814. doi: 10.4103/1673-5374.206656

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Colcombe, S., and Kramer, A. F. (2003). Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol. Sci.* 14, 125–130. doi: 10.1111/1467-9280.t01-1-01430

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Colcombe, S. J., Erickson, K. I., Scalf, P. E., Kim, J. S., Prakash, R., McAuley, E., et al. (2006). Aerobic exercise training increases brain volume in aging humans. *J. Gerontol. A Biol. Sci. Med. Sci.* 61, 1166–1170. doi: 10.1093/gerona/61.11.1166

[CrossRef Full Text](#) | [Google Scholar](#)

Conti, M. Z., Vicini-Chilovi, B., Riva, M., Zanetti, M., Liberini, P., Padovani, A., et al. (2013). Odor identification deficit predicts clinical conversion from mild cognitive impairment to dementia due to Alzheimer's disease. *Arch. Clin. Neuropsychol.* 28, 391–399. doi: 10.1093/arclin/act032

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Cosetti, M. K., Pinkston, J. B., Flores, J. M., Friedmann, D. R., Jones, C. B., Roland, J. T. Jr., et al. (2016). Neurocognitive testing and cochlear implantation: insights into performance in older adults. *Clin. Intervent. Aging* 11, 603–613. doi: 10.2147/CIA.S100255

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Costa, D. A., Cracchiolo, J. R., Bachstetter, A. D., Hughes, T. F., Bales, K. R., Paul, S. M., et al. (2007). Enrichment improves cognition in AD mice by amyloid-related and unrelated mechanisms. *Neurobiol. Aging* 28, 831–844. doi: 10.1016/j.neurobiolaging.2006.04.009

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Costello, E., Kafchinski, M., Vrazel, J., and Sullivan, P. (2011). Motivators, barriers, and beliefs regarding physical activity in an older adult population. *J. Geriatr. Phys. Ther.* 34, 138–147. doi: 10.1519/JPT.0b013e31820e0e71

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Cotman, C. W., Berchtold, N. C., and Christie, L. A. (2007). Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends Neurosci.* 30, 464–472. doi: 10.1016/j.tins.2007.06.011

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Damm, M., Pikart, L. K., Reimann, H., Burkert, S., Göktas, Ö., Haxel, B., et al. (2014). Olfactory training is helpful in postinfectious olfactory loss: a randomized, controlled, multicenter study. *Laryngoscope* 124, 826–831. doi: 10.1002/lary.24340

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Dawes, P., Emsley, R., Cruickshanks, K. J., Moore, D. R., Fortnum, H., Edmondson-Jones, M., et al. (2015). Hearing loss and cognition: the role of hearing aids, social isolation and depression. *PLoS One* 10: e0119616. doi: 10.1371/journal.pone.0119616

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

De Cicco, V., Barresi, M., Tramonti Fantozzi, M. P., Cataldo, E., Parisi, V., and Manzoni, D. (2016). Oral implant-prostheses: new teeth for a brighter brain. *PLoS One* 11: e0148715. doi: 10.1371/journal.pone.0148715

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Del Brutto, O. H., Gardener, H., Del Brutto, V. J., Maestre, G. E., Zambrano, M., Montenegro, J. E., et al. (2014). Edentulism associates with worse cognitive performance in community-dwelling elders in rural Ecuador: results of the Atahualpa project. *J. Commun. Health* 39, 1097–1100. doi: 10.1007/s10900-014-9857-3

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Devanand, D. P., Michaels-Marston, K. S., Liu, X., Pelton, G. H., Padilla, M., Marder, K., et al. (2000). Olfactory deficits in patients with mild cognitive impairment predict Alzheimer's disease at follow-up. *Am. J. Psychiatry* 157, 1399–1405. doi: 10.1176/appi.ajp.157.9.1399

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Devanand, D. P., Tabert, M. H., Cuasay, K., Manly, J. J., Schupf, N., Brickman, A. M., et al. (2010). Olfactory identification deficits and MCI in a multi-ethnic elderly community sample. *Neurobiol. Aging* 31, 1593–1600. doi: 10.1016/j.neurobiolaging.2008.09.008

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Dickinson, C., and Rabbitt, P. (1991). Simulated visual impairment: effects on text comprehension and reading speed. *Clin. Vis. Sci.* 6, 301–308.

[Google Scholar](#)

Dong, J., Pinto, J. M., Guo, X., Alonso, A., Tranah, G., Cauley, J. A., et al. (2017). The prevalence of anosmia and associated factors among U. S. black and white older adults. *J. Gerontol. A Biol. Sci. Med. Sci.* 72, 1080–1086. doi: 10.1093/gerona/glx081

[CrossRef Full Text](#) | [Google Scholar](#)

Doty, R. L., Deems, D. A., and Stellar, S. (1988). Olfactory dysfunction in Parkinsonism: a general deficit unrelated to neurologic signs, disease stage, or disease duration. *Neurology* 38, 1237–1244. doi: 10.1212/WNL.38.8.1237

[CrossRef Full Text](#) | [Google Scholar](#)

Doty, R. L., Shaman, P., Applebaum, S. L., Giberson, R., Siksorski, L., and Rosenberg, L. (2015). Smell identification ability: changes with age. *Science* 226, 1441–1443. doi: 10.1126/science.6505700

[CrossRef Full Text](#) | [Google Scholar](#)

Downs, J., Rodger, J., Li, C., Tan, X., Hu, N., Wong, K., et al. (2018).

Environmental enrichment intervention for Rett syndrome: an individually randomised stepped wedge trial. *Orphanet J. Rare Dis.* 13: 3. doi: 10.1186/s13023-017-0752-8

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Drobny, J. V., Anstey, K. J., and Andrews, S. (2005). Visual memory testing in older adults with age-related visual decline: a measure of memory performance or visual functioning? *J. Clin. Exp. Neuropsychol.* 27, 425–435. doi: 10.1080/138033990520241

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Eckert, M. A., Cute, S. L., Vaden, K. I. Jr., Kuchinsky, S. E., and Dubno, J. R. (2012). Auditory cortex signs of age-related hearing loss. *J. Assoc. Res. Otolaryngol.* 13, 703–713. doi: 10.1007/s10162-012-0332-5

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Eckert, M. A., Walczak, A., Ahlstrom, J., Denslow, S., Horwitz, A., and Dubno, J. R. (2008). Age-related effects on word recognition: reliance on cognitive control systems with structural declines in speech-responsive cortex. *J. Assoc. Res. Otolaryngol.* 9, 252–259. doi: 10.1007/s10162-008-0113-3

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Elliott, A. F., McGwin, G. Jr., and Owsley, C. (2009). Vision-enhancing interventions in nursing home residents and their short-term effect on physical and cognitive function. *J. Am. Geriatr. Soc.* 57, 202–208. doi: 10.1111/j.1532-5415.2008.02099.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Elsig, F., Schimmel, M., Duvernay, E., Giannelli, S. V., Graf, C. E., Carlier, S., et al. (2015). Tooth loss, chewing efficiency and cognitive impairment in geriatric patients. *Gerontology* 32, 149–156. doi: 10.1111/ger.12079

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Erb, J., Henry, M. J., Eisner, F., and Obleser, J. (2013). The brain dynamics of rapid perceptual adaptation to adverse listening conditions. *J. Neurosci.* 33, 10688–10697. doi: 10.1523/JNEUROSCI.4596-12.2013

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Erb, J., and Obleser, J. (2013). Upregulation of cognitive control networks in older adults' speech comprehension. *Front. Syst. Neurosci.* 7: 116. doi: 10.3389/fnsys.2013.00116

[CrossRef Full Text](#) | [Google Scholar](#)

Erickson, K. I., Gildengers, A. G., and Butters, M. A. (2013). Physical activity and brain plasticity in late adulthood. *Dialog. Clin. Neurosci.* 15, 99–108.

[Google Scholar](#)

Erickson, K. I., Leckie, R. L., and Weinstein, A. M. (2014). Physical activity, fitness, and gray matter volume. *Neurobiol. Aging* 35(Suppl. 2), S20–S28. doi: 10.1016/j.neurobiolaging.2014.03.034

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Erickson, K. I., Voss, M. W., Prakash, R. S., Basak, C., Szabo, A., Chaddock, L., et al. (2011). Exercise training increases size of hippocampus and improves memory. *Proc. Natl. Acad. Sci. U. S. A.* 108, 3017–3022. doi: 10.1073/pnas.1015950108

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ferris, L. T., Williams, J. S., and Shen, C. L. (2007). The effect of acute exercise on serum brain-derived neurotrophic factor levels and cognitive function. *Med. Sci. Sports Exerc.* 39, 728–734. doi: 10.1249/mss.0b013e31802f04c7

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Fukushima-Nakayama, Y., Ono, T., Hayashi, M., Inoue, M., Wake, H., Ono, T., et al. (2017). Reduced mastication impairs memory function. *J. Dent. Res.* 96, 1058–1066. doi: 10.1177/0022034517708771

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Geißler, K., Reimann, H., Gudziol, H., Bitter, T., and Guntinas-Lichius, O. (2014). Olfactory training for patients with olfactory loss after upper respiratory tract infections. *Eur. Arch. Otorhinolaryngol.* 271, 1557–1562. doi: 10.1007/s00405-013-2747-y

<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Gil-Montoya, J. A., Barrios, R., Santana, S., Sanchez-Lara, I., Pardo, C. C., Fornieles-Rubio, F., et al. (2015). Is periodontitis a risk factor for cognitive impairment and dementia? A case-control study. *J. Periodont.* 86, 244–253. doi: 10.1902/jop.2014.140340

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Gopinath, B., McMahon, C. M., Burlutsky, G., and Mitchell, P. (2016). Hearing and vision impairment and the 5-year incidence of falls in older adults. *Age Ageing* 45, 409–414. doi: 10.1093/ageing/afw022

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Gopinath, B., Schneider, J., McMahon, C. M., Burlutsky, G., Leeder, S. R., and Mitchell, P. (2013). Dual sensory impairment in older adults increases the risk of mortality: a population-based study. *PLoS One* 8: e55054. doi: 10.1371/journal.pone.0055054

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Gothe, N. P., and Kendall, B. J. (2016). Barriers, motivations, and preferences for physical activity among female African American older adults. *Gerontol. Geriatr. Med.* 2: 2333721416677399. doi: 10.1177/2333721416677399

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Grabe, H. J., Schwahn, C., Völzke, H., Spitzer, C., Freyberger, H. J., John, U., et al. (2009). Tooth loss and cognitive impairment. *J. Clin. Periodontol.* 36, 550–557. doi: 10.1111/j.1600-051X.2009.01426.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Gregory, M. A., Boa Sorte Silva, N. C., Gill, D. P., McGowan, C. L., Liu-Ambrose, T., Shoemaker, J. K., et al. (2017). Combined dual-task gait training and aerobic exercise to improve cognition, mobility, and vascular health in community-dwelling older adults at risk for future cognitive decline. *J. Alzheimers Dis.* 57, 747–763. doi: 10.3233/JAD-161240

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Griffin, S. O., Jones, J. A., Brunson, D., Griffin, P. M., and Bailey, W. D. (2012). Burden of oral disease among older adults and implications for public health priorities. *Am. J. Public Health* 102, 411–418. doi: 10.2105/AJPH.2011.300362

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Haehner, A., Tosch, C., Wolz, M., Klingelhoefer, L., Fauser, M., Storch, A., et al. (2013). Olfactory training in patients with Parkinson's disease. *PLoS One* 8: e61680. doi: 10.1371/journal.pone.0061680

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Håkansson, K., Ledreux, A., Daffner, K., Terjestam, Y., Bergman, P., Carlsson, R., et al. (2017). BDNF responses in healthy older persons to 35 minutes of physical exercise, cognitive training, and mindfulness: associations with <https://assignbuster.com/environmental-enrichment-and-successful-aging/>

working memory function. *J. Alzheimers Dis.* 55, 645–657. doi: 10. 3233/JAD-160593

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hanna-Pladdy, B., and MacKay, A. (2011). The relation between instrumental musical activity and cognitive aging. *Neuropsychology* 25, 378–386. doi: 10. 1037/a0021895

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Harris, K. C., Dubno, J. R., Keren, N. I., Ahlstrom, J. B., and Eckert, M. A. (2009). Speech recognition in younger and older adults: a dependency on low-level auditory cortex. *J. Neurosci.* 29, 6078–6087. doi: 10. 1523/JNEUROSCI. 0412-09. 2009

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Heine, C., and Browning, C. J. (2002). Communication and psychosocial consequences of sensory loss in older adults: overview and rehabilitation directions. *Disabil. Rehabil.* 24, 763–773. doi: 10. 1080/09638280210129162

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hernowo, A. T., Prins, D., Baseler, H. A., Plank, T., Gouws, A. D., Hooymans, J. M., et al. (2014). Morphometric analyses of the visual pathways in macular degeneration. *Cortex* 56, 99–110. doi: 10. 1016/j.cortex. 2013. 01. 003

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hertzog, C., Kramer, A. F., Wilson, R. S., and Lindenberger, U. (2008). Enrichment effects on adult cognitive development: can the functional capacity of older adults be preserved and enhanced? *Psychol. Sci. Public Interest* 9, 1–65. doi: 10.1111/j.1539-6053.2009.01034.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hirano, Y., Obata, T., Takahashi, H., Tachibana, A., Kuroiwa, D., Takahashi, T., et al. (2013). Effects of chewing on cognitive processing speed. *Brain Cogn.* 81, 376–381. doi: 10.1016/j.bandc.2012.12.002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hirano, Y., and Onozuka, M. (2015). Chewing and attention: a positive effect on sustained attention. *Biomed Res. Int.* 2015: 367026. doi: 10.1155/2015/367026

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hoffman, H. J., Ishii, E. K., and MacTurk, R. H. (1998). Age-related changes in the prevalence of smell/taste problems among the United States adult population. Results of the 1994 disability supplement to the National Health Interview Survey (NHIS). *Ann. N. Y. Acad. Sci.* 855, 716–722. doi: 10.1111/j.1749-6632.1998.tb10650.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hoffman, H. J., Rawal, S., Li, C.-M., and Duffy, V. B. (2016). New chemosensory component in the U. S. National Health and Nutrition Examination Survey (NHANES): first-year results for measured olfactory  
<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

dysfunction. *Rev. Endocr. Metab. Disord.* 17, 221–240. doi: 10. 1007/s11154-016-9364-1

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Humes, L. E., Busey, T. A., Craig, J., and Kewley-Port, D. (2013). Are age-related changes in cognitive function driven by age-related changes in sensory processing? *Atten. Percept. Psychophys.* 75, 508–524. doi: 10.3758/s13414-012-0406-9

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Hummel, T., Rissom, K., Reden, J., Hähner, A., Weidenbecher, M., and Hüttenbrink, K.-B. (2009). Effects of olfactory training in patients with olfactory loss. *Laryngoscope* 119, 496–499. doi: 10. 1002/lary. 20101

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Husain, F. T., Medina, R. E., Davis, C. W., Szymko-Bennett, Y., Simonyan, K., Pajor, N. M., et al. (2011). Neuroanatomical changes due to hearing loss and chronic tinnitus: a combined VBM and DTI study. *Brain Res.* 1369, 74–88. doi: 10. 1016/j. brainres. 2010. 10. 095

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ichihashi, Y., Saito, N., Arakawa, Y., Kurata, C., Iinuma, M., Tamura, Y., et al. (2008). The bite-raised condition in aged SAMP8 mice reduces the expression of glucocorticoid receptors in the dorsal and ventral hippocampus. *Okajimas Folia Anat. Jpn.* 84, 137–142. doi: 10. 2535/ofaj. 84. 137

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ishii, K., Kabata, T., and Oshika, T. (2008). The impact of cataract surgery on cognitive impairment and depressive mental status in elderly patients. *Am. J. Ophthalmol.* 146, 404–409. doi: 10.1016/j.ajo.2008.05.014

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Jankowsky, J. L., Melnikova, T., Fadale, D. J., Xu, G. M., Slunt, H. H., Gonzales, V., et al. (2005). Environmental enrichment mitigates cognitive deficits in a mouse model of Alzheimer's disease. *J. Neurosci.* 25, 5217–5224. doi: 10.1523/JNEUROSCI.5080-04.2005

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Jefferis, J. M., Clarke, M. P., and Taylor, J. P. (2015). Effect of cataract surgery on cognition, mood, and visual hallucinations in older adults. *J. Cataract Refract. Surg.* 41, 1241–1247. doi: 10.1016/j.jcrs.2014.09.044

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Jonasson, L. S., Nyberg, L., Kramer, A. F., Lundquist, A., Riklund, K., and Boraxbekk, C. J. (2017). Aerobic exercise intervention, cognitive performance, and brain structure: results from the Physical Influences on Brain in Aging (PHIBRA) Study. *Front. Aging Neurosci.* 8: 336. doi: 10.3389/fnagi.2016.00336

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kamer, A. R., Morse, D. E., Holm-Pedersen, P., Mortensen, E. L., and Avlund, K. (2012). Periodontal inflammation in relation to cognitive function in an older adult Danish population. *J. Alzheimers Dis.* 28, 613–624. doi: 10.3233/JAD-2011-102004

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kamiya, K., Narita, N., and Iwaki, S. (2016). Improved prefrontal activity and chewing performance as function of wearing denture in partially edentulous elderly individuals: functional near-infrared spectroscopy study. *PLoS One* 11: e0158070. doi: 10.1371/journal.pone.0158070

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kato, T., Usami, T., Noda, Y., Hasegawa, M., Ueda, M., and Nabeshima, T. (1997). The effect of the loss of molar teeth on spatial memory and acetylcholine release from the parietal cortex in aged rats. *Behav. Brain Res.* 83, 239–242. doi: 10.1016/S0166-4328(97)86078-0

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Keller, B. K., Morton, J. L., Thomas, V. S., and Potter, J. F. (1999). The effect of visual and hearing impairments on functional status. *J. Am. Geriatr. Soc.* 47, 1319–1325. doi: 10.1111/j.1532-5415.1999.tb07432.x

[CrossRef Full Text](#) | [Google Scholar](#)

Kemoun, G., Thibaud, M., Roumagne, N., Carette, P., Albinet, C., Toussaint, L., et al. (2010). Effects of a physical training programme on cognitive

function and walking efficiency in elderly persons with dementia. *Dement. Geriatr. Cogn. Disord.* 29, 109–114. doi: 10. 1159/000272435

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kennedy, G., Hardman, R. J., Macpherson, H., Scholey, A. B., and Pipingas, A. (2017). How does exercise reduce the rate of age-associated cognitive decline? A review of potential mechanisms. *J. Alzheimers Dis.* 55, 1–18. doi: 10. 3233/JAD-160665

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kimoto, K., Ono, Y., Tachibana, A., Hirano, Y., Otsuka, T., Ohno, A., et al. (2011). Chewing-induced regional brain activity in edentulous patients who received mandibular implant-supported overdentures: a preliminary report. *J. Prosthodont. Res.* 55, 89–97. doi: 10. 1016/j.jpor. 2010. 09. 006

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kimura, Y., Ogawa, H., Yoshihara, A., Yamaga, T., Takiguchi, T., Wada, T., et al. (2013). Evaluation of chewing ability and its relationship with activities of daily living, depression, cognitive status and food intake in the community-dwelling elderly. *Geriatr. Gerontol. Int.* 13, 718–725. doi: 10. 1111/ggi. 12006

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kirk-Sanchez, N. J., and McGough, E. L. (2014). Physical exercise and cognitive performance in the elderly: current perspectives. *Clin. Interv. Aging* 9, 51–62. doi: 10. 2147/CIA. S39506

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Klaver, C. C., Wolfs, R. C., Vingerling, J. R., Hofman, A., and de Jong, P. T. (1998). Age-specific prevalence and causes of blindness and visual impairment in an older population: the Rotterdam Study. *Arch. Ophthalmol.* 116, 653–658. doi: 10.1001/archopht.116.5.653

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kohli, P., Soler, Z. M., Nguyen, S. A., Muus, J. S., and Schlosser, R. J. (2016). The association between olfaction and depression: a systematic review. *Chem. Senses* 41, 479–486. doi: 10.1093/chemse/bjw061

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kollendorfer, K., Jakab, A., Mueller, C., Trattnig, S., and Schopf, V. (2015). Effects of chronic peripheral olfactory loss on functional brain networks. *Neuroscience* 310, 589–599. doi: 10.1016/j.neuroscience.2015.09.045

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Konstantinidis, I., Tsakiroglou, E., Bekiaridou, P., Kazantzidou, C., and Constantinidis, J. (2013). Use of olfactory training in post-traumatic and postinfectious olfactory dysfunction. *Laryngoscope* 123, E85–E90. doi: 10.1002/lary.24390

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Konstantinidis, I., Tsakiroglou, E., and Constantinidis, J. (2016). Long term effects of olfactory training in patients with postinfectious olfactory loss. *Laryngoscope* 54, 170–175. doi: 10. 4193/Rhin15. 264

[PubMed Abstract](#) | [CrossRef Full Text](#)

Kubo, K., Chen, H., Zhou, X., Liu, J., and Darbin, O. (2015). Chewing, stress-related diseases, and brain function. *Biomed Res. Int.* 2015: 412493. doi: 10. 1155/2015/412493

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Kubo, K. Y., Yamada, Y., Iinuma, M., Iwaku, F., Tamura, Y., Watanabe, K., et al. (2007). Occlusal disharmony induces spatial memory impairment and hippocampal neuron degeneration via stress in SAMP8 mice. *Neurosci. Lett.* 414, 188–191. doi: 10. 1016/j. neulet. 2006. 12. 020

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lafaille-Magnan, M.-E., Poirier, J., Etienne, P., Tremblay-Mercier, J., Frenette, J., Rosa-Neto, P., et al. (2017). Odor identification as a biomarker of preclinical AD in older adults at risk. *Neurology* 89, 327–335. doi: 10. 1212/WNL. 0000000000004159

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lai, S. W., Lin, C. L., and Liao, K. F. (2014). Cataract may be a non-memory feather of Alzheimer's disease in older people. *Eur. J. Epidemiol.* 29, 405–409. doi: 10. 1007/s10654-014-9903-6

[PubMed Abstract](#) | [CrossRef Full Text](#)

Lazarov, O., Robinson, J., Tang, Y. P., Hairston, I. S., Korade-Mirnics, Z., Lee, V. M., et al. (2005). Environmental enrichment reduces Abeta levels and amyloid deposition in transgenic mice. *Cell* 120, 701–713. doi: 10.1016/j.cell.2005.01.015

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Leporé, N., Shi, Y., Lepore, F., Fortin, M., Voss, P., Chou, Y. Y., et al. (2009). Patterns of hippocampal shape and volume differences in blind subjects. *Neuroimage* 46, 949–957. doi: 10.1016/j.neuroimage.2009.01.071

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lexomboon, D., Trulsson, M., Wård, I., and Parker, M. G. (2012). Chewing ability and tooth loss: association with cognitive impairment in an elderly population study. *J. Am. Geriatr. Soc.* 60, 1951–1956. doi: 10.1111/j.1532-5415.2012.04154.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Li, W., Howard, J. D., and Gottfried, J. A. (2010). Disruption of odour quality coding in piriform cortex mediates olfactory deficits in Alzheimer's disease. *Brain* 133, 2714–2726. doi: 10.1093/brain/awq209

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, F. R., Chien, W. W., Li, L., Niparko, J. K., and Francis, H. W. (2012). Cochlear implantation in older adults. *Medicine* 91, 229–241. doi: 10.1097/MD.0b013e31826b145a

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, F. R., Ferrucci, L., An, Y., Goh, J. O., Doshi, J., Metter, E. J., et al. (2014). Association of hearing impairment with brain volume changes in older adults. *Neuroimage* 90, 84–92. doi: 10.1016/j.neuroimage.2013.12.059

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, F. R., Metter, E. J., O'Brien, R. J., Resnick, S. M., Zonderman, A. B., and Ferrucci, L. (2011a). Hearing loss and incident dementia. *Arch. Neurol.* 68, 214–220. doi: 10.1001/archneurol.2010.362

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, F. R., Thorpe, R., Gordon-Salant, S., and Ferrucci, L. (2011b). Hearing loss prevalence and risk factors among older adults in the United States. *J. Gerontol. A Biol. Sci. Med. Sci.* 66A, 582–590. doi: 10.1093/gerona/glr002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, H., Zhang, L., Chen, W., Zhu, Y., Chen, C., Chan, K. C., et al. (2018). Visual restoration after cataract surgery promotes functional and structural brain recovery. *EBioMedicine* doi: 10.1016/j.ebiom.2018.03.002 [Epub ahead of print].

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lin, M. Y., Gutierrez, P. R., Stone, K. L., Yaffe, K., Ensrud, K. E., Fink, H. A., et al. (2004). Vision impairment and combined vision and hearing impairment predict cognitive and functional decline in older women. *J. Am. Geriatr. Soc.* 52, 1996–2002. doi: 10.1111/j.1532-5415.2004.52554.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lindenberger, U., and Baltes, P. B. (1994). Sensory functioning and intelligence in old age: a strong connection. *Psychol. Aging* 9, 339–355. doi: 10.1037/0882-7974.9.3.339

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lipardo, D. S., Aseron, A. M. C., Kwan, M. M., and Tsang, W. W. (2017). Effect of exercise and cognitive training on falls and fall-related factors in older adults with mild cognitive impairment: a systematic review. *Arch. Phys. Med. Rehabil.* 98, 2079–2096. doi: 10.1016/j.apmr.2017.04.021

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Liu, G., Zong, G., Doty, R. L., and Sun, Q. (2016). Prevalence and risk factors of taste and smell impairment in a nationwide representative sample of the US population: a cross-sectional study. *BMJ Open* 6: e013246. doi: 10.1136/bmjopen-2016-013246

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Lou, A. R., Madsen, K. H., Julian, H. O., Toft, P. B., Kjaer, T. W., Paulson, O. B., et al. (2013). Postoperative increase in grey matter volume in visual cortex

after unilateral cataract surgery. *Acta Ophthalmol.* 91, 58–65. doi: 10. 1111/j. 1755-3768. 2011. 02304. x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Loughrey, D. G., Kelly, M. E., Kelley, G. A., Brennan, S., and Lawlor, B. A. (2018). Association of age-related hearing loss with cognitive function, cognitive impairment, and dementia: a systematic review and meta-analysis. *JAMA Otolaryngol. Head Neck Surg.* 144, 115–126. doi: 10. 1001/jamaoto. 2017. 2513

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Luo, J., Wu, B., Zhao, Q., Guo, Q., Meng, H., Yu, L., et al. (2015). Association between tooth loss and cognitive function among 3063 Chinese older adults: a community-based study. *PLoS One* 10: e0120986. doi: 10. 1371/journal. pone. 0120986

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Luzzi, S., Snowden, J. S., Neary, D., Coccia, M., Provinciali, L., and Lambon Ralph, M. A. (2007). Distinct patterns of olfactory impairment in Alzheimer's disease, semantic dementia, frontotemporal dementia, and corticobasal degeneration. *Neuropsychologia* 45, 1823–1831. doi: 10. 1016/j. neuropsychologia. 2006. 12. 008

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Matsuyama, Y., Aida, J., Ysuboya, T., Hikichi, H., Kondo, K., Kawachi, I., et al. (2017). Are lowered socioeconomic circumstances casually related to tooth <https://assignbuster.com/environmental-enrichment-and-successful-aging/>

loss? A natural experiment involving the 2011 Great East Japan earthquake. *Am. J. Epidemiol.* 186, 54–62. doi: 10. 1093/aje/kwx059

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mavros, Y., Gates, N., Wilson, G. C., Jain, N., Meiklejohn, J., Brodaty, H., et al. (2017). Mediation of cognitive function improvements by strength gains after resistance training in older adults with mild cognitive impairment: outcomes of the study of mental and resistance training. *J. Am. Geriatr. Soc.* 65, 550–559. doi: 10. 1111/jgs. 14542

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

McGregor, K. M., Crosson, B., Mammino, K., Omar, J., Garcia, P. S., and Nocera, J. R. (2018). Influences of 12-week physical activity interventions on TMS measures of cortical network inhibition and upper extremity motor performance in older adults- A feasibility study. *Front. Aging Neurosci.* 9: 422. doi: 10. 3389/fnagi. 2017. 00422

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Meijer, E. P., Goris, A. H., van Dongen, J. L., Bast, A., and Westerterp, K. R. (2002). Exercise-induced oxidative stress in older adults as a function of habitual activity level. *J Am. Geriatr. Soc.* 50, 349–353. doi: 10. 1046/j. 1532-5415. 2002. 50069. x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Meusel, T., Westermann, B., Fuhr, P., Hummel, T., and Welge-Lüssen, A. (2010). The course of olfactory deficits in patients with Parkinson's disease - <https://assignbuster.com/environmental-enrichment-and-successful-aging/>

a study based on psychophysical and electrophysiological measures.

*Neurosci. Lett.* 486, 166–170. doi: 10.1016/j.neulet.2010.09.044

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mick, P., Kawachi, I., and Lin, F. R. (2014). The association between hearing loss and social isolation in older adults. *Otolaryngol. Head Neck Surg.* 150, 378–384. doi: 10.1177/0194599813518021

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mine, M., Miyata, K., Morikawa, M., Nishi, T., Okamoto, N., Kawasaki, R., et al. (2016). Association of visual acuity and cognitive impairment in older individuals: Fujiwara-kyo eye study. *Biores. Open Access* 5, 228–234. doi: 10.1089/biores.2016.0023

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mitoku, K., Masaki, N., Ogata, Y., and Okamoto, K. (2016). Vision and hearing impairments, cognitive impairment and mortality among long-term care recipients: a population-based cohort study. *BMC Geriatr.* 16: 112. doi: 10.1186/s12877-016-0286-2

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Miyake, S., Yoshikawa, G., Yamada, K., Sasaguri, K., Yamamoto, T., Onozuka, M., et al. (2012). Chewing ameliorates stress-induced suppression of spatial memory by increasing glucocorticoid receptor expression in the hippocampus. *Brain Res.* 1446, 34–39. doi: 10.1016/j.brainres.2012.01.011

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Miyata, K., Obayashi, K., Saeki, K., Tone, N., Tanaka, K., Nishi, T., et al. (2016). Higher cognitive function in elderly individuals with previous cataract surgery: cross-sectional association independent of visual acuity in the HEIJO-KYO cohort. *Rejuvenation Res.* 19, 239–243. doi: 10.1089/rej.2015.1718

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mortimer, J. A., Ding, D., Borenstein, A. R., DeCarli, C., Guo, Q., Wu, Y., et al. (2012). Changes in brain volume and cognition in a randomized trial of exercise and social interaction in a community-based sample of non-demented Chinese elders. *J. Alzheimers Dis.* 30, 757–766. doi: 10.3233/JAD-2012-120079

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mosnier, I., Bebear, J. P., Marx, M., Fraysse, B., Truy, E., Lina-Granade, G., et al. (2015). Improvement of cognitive function after cochlear implantation in elderly patients. *JAMA Otolaryngol. Head Neck Surg.* 141, 442–450. doi: 10.1001/jamaoto.2015.129

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Mulrow, C. D., Aguilar, C., Endicott, J. E., Tuley, M. R., Velez, R., Charlip, W. S., et al. (1990). Quality-of-life changes and hearing impairment: a randomized trial. *Ann. Intern. Med.* 113, 188–194. doi: 10.7326/0003-4819-113-3-188

[CrossRef Full Text](#) | [Google Scholar](#)

Mummolo, S., Ortù, E., Necozione, S., Monaco, A., and Marzo, G. (2014).

Relationship between mastication and cognitive function in elderly in L'Aquila. *Int. J. Clin. Exp. Med.* 7, 1040–1046.

[Google Scholar](#)

Muñoz, B., West, S. K., Rubin, G. S., Schein, O. D., Quigley, H. A., Bressler, S. B., et al. (2000). Causes of blindness and visual impairment in a population of older Americans: the Salisbury eye evaluation study. *Arch. Ophthalmol.* 118, 819–825. doi: 10.1001/archopht.118.6.819

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Murphy, C., Schubert, C. R., Cruickshanks, K. J., Klein, B. E., Klein, R., and Nondahl, D. M. (2002). Prevalence of olfactory impairment in older adults. *JAMA* 288, 2307–2312. doi: 10.1001/jama.288.18.2307

[CrossRef Full Text](#) | [Google Scholar](#)

Narita, N., Kamiya, K., Yamamura, K., Kawasaki, S., Matsumoto, T., and Tanaka, N. (2009). Chewing-related prefrontal cortex activation while wearing partial denture prosthesis: pilot study. *J. Prosthodont. Res.* 53, 126–135. doi: 10.1016/j.jpor.2009.02.005

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Nguyen, A. D., Shenton, M. E., and Levitt, J. J. (2010). Olfactory dysfunction in schizophrenia: a review of neuroanatomy and psychophysiological

measurements. *Harv. Rev. Psychiatry* 18, 279–292. doi: 10. 1016/j.bbr. 2008. 07. 008

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Niemann, C., Godde, B., and Voelcker-Rehage, C. (2014). Not only cardiovascular, but also coordinative exercise increases hippocampal volume in older adults. *Front. Aging Neurosci.* 6: 170. doi: 10. 3389/fnagi. 2014. 00170

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Noble, J. M., Borrell, L. N., Papapanou, P. N., Elkind, M. S., Scarmeas, N., and Wright, C. B. (2009). Periodontitis is associated with cognitive impairment among older adults: analysis of NHANES-III. *J. Neurol. Neurosurg. Psychiatry* 80, 1206–1211. doi: 10. 1136/jnnp. 2009. 174029

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Nocera, J., Crosson, B., Mammino, K., and McGregor, K. M. (2017). Changes in cortical activation patterns in language areas following an aerobic exercise intervention in older adults. *Neural Plast.* 2017: 6340302. doi: 10. 1155/2017/6340302

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Norhey, J. M., Cherbuin, N., Pumpa, K. L., Smee, D. J., and Rattray, B. (2018). Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br. J. Sports Med.* 52, 154–160. doi: 10. 1136/bjsports-2016-096587

<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Okamoto, N., Morikawa, M., Okamoto, K., Habu, N., Hazaki, K., Harano, A., et al. (2010). Tooth loss is associated with mild memory impairment in the elderly: the Fujiwara-kyo study. *Brain Res.* 1349, 68–75. doi: 10.1016/j.brainres.2010.06.054

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Okihara, H., Ito, J., Kokai, S., Ishida, T., Hiranuma, M., Kato, C., et al. (2014). Liquid diet induces memory impairment accompanied by a decreased number of hippocampal neurons in mice. *J. Neurosci. Res.* 92, 1010–1017. doi: 10.1002/jnr.23383

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Olze, H., Gräbel, S., Förster, U., Zirke, N., Huhnd, L. E., Haupt, H., et al. (2012). Elderly patients benefit from cochlear implantation regarding auditory rehabilitation, quality of life, tinnitus, and stress. *Laryngoscope* 122, 196–203. doi: 10.1002/lary.22356

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ong, S. Y., Cheung, C. Y., Li, X., Lamoureux, E. L., Ikram, M. K., Ding, J., et al. (2012). Visual impairment, age-related eye diseases, and cognitive function: the Singapore Malay Eye Study. *Arch. Ophthalmol.* 130, 895–900. doi: 10.1001/archophthalmol.2012.152

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ono, Y., Yamamoto, T., Kubo, K. Y., and Onozuka, M. (2010). Occlusion and brain function: mastication as a prevention of cognitive dysfunction. *J. Oral Rehabil.* 37, 624–640. doi: 10. 1111/j. 1365-2842. 2010. 02079. x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Onozuka, M., Fujita, M., Watanabe, K., Hirano, Y., Niwa, M., Nishiyama, K., et al. (2002). Mapping brain region activity during chewing: a functional magnetic resonance imaging study. *J. Dent. Res.* 81, 743–746. doi: 10. 1177/0810743

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Onozuka, M., Watanabe, K., Mirbod, S. M., Ozono, S., Nishiyama, K., Karasawa, N., et al. (1999). Reduced mastication stimulates impairment of spatial memory and degeneration of hippocampal neurons in aged SAMP8 mice. *Brain Res.* 826, 148–153. doi: 10. 1016/S0006-8993(99)01255-X

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Parbery-Clark, A., Anderson, S., Hittner, E., and Kraus, N. (2012). Musical experience offsets age-related delays in neural timing. *Neurobiol. Aging* 33, 1483. e1–1483. e4. doi: 10. 1016/j. neurobiolaging. 2011. 12. 015

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Parrao, T., Chana, P., Venegas, P., Behrens, M. I., and Aylwin, M. L. (2012). Olfactory deficits and cognitive dysfunction in Parkinson's disease. *Neurodegener. Dis.* 10, 179–182. doi: 10. 1159/000335915

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Patel, T. R. (2012). Environmental enrichment: aging and memory. *Yale J. Biol. Med.* 85, 491–500.

[Google Scholar](#)

Patel, Z. M., Wise, S. K., and DelGaudio, J. M. (2017). Randomized controlled trial demonstrating cost-effective method of olfactory training in clinical practice: essential oils at uncontrolled concentration. *Laryngoscope Investig. Otolaryngol.* 2, 53–56. doi: 10.1002/lio2.62

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Peelle, J. E., Troiani, V., Grossman, M., and Wingfield, A. (2011). Hearing loss in older adults affects neural systems supporting speech comprehension. *J. Neurosci.* 31, 12638–12643. doi: 10.1523/JNEUROSCI.2559-11.2011

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Pekala, K., Chandra, R. K., and Turner, J. H. (2016). Efficacy of olfactory training in patients with olfactory loss: a systematic review and meta-analysis. *Int. Forum Allergy Rhinol.* 6, 299–307. doi: 10.1002/alr.21669

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Peng, P., Gu, H., Xiao, W., Si, L. F., Wang, J. F., Wang, S. K., et al. (2013). A voxel-based morphometry study of anosmic patients. *Br. J. Radiol.* 86: 20130207. doi: 10.1259/bjr.20130207

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

<https://assignbuster.com/environmental-enrichment-and-successful-aging/>

Pereira, A. C., Huddleston, D. E., Brickman, A. M., Sosunov, A. A., Hen, R., McKhann, G. M., et al. (2007). An in vivo correlate of exercise-induced neurogenesis in the adult dentate gyrus. *Proc. Natl. Acad. Sci. U. S. A.* 104, 5638–5643. doi: 10.1073/pnas.0611721104

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Peres, M. A., Bastos, J. L., Watt, R. G., Xavier, A. J., Barbato, P. R., and D'Orsi, E. (2014). Tooth loss is associated with severe cognitive impairment among older people: findings from a population-based study in Brazil. *Aging Ment. Health* 19, 876–884. doi: 10.1080/13607863.2014.977770

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Peters, J. M., Hummel, T., Kratzsch, T., Lötsch, J., Skarke, C., and Frölich, L. (2003). Olfactory function in mild cognitive impairment and Alzheimer's disease: an investigation using psychophysical and electrophysiological techniques. *Am. J. Psychiatry* 160, 1995–2002. doi: 10.1176/appi.ajp.160.11.1995

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Pichora-Fuller, E. (2006). Perceptual effort and apparent cognitive decline: implications for audiology rehabilitation. *Semin. Hear.* 27, 284–293. doi: 10.1055/s-2006-954855

[CrossRef Full Text](#) | [Google Scholar](#)

Polito, L., Chierchia, A., Tunisi, M., Bouybayoune, I., Kehoe, P. G., Albani, P., et al. (2014). Environmental enrichment lessens cognitive decline in app23. <https://assignbuster.com/environmental-enrichment-and-successful-aging/>

mice without affecting brain sirtuin expression. *J. Alzheimers Dis.* 42, 851–864. doi: 10. 3233/JAD-131430

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ponsen, M. M., Stoffers, D., Booij, J., van Eck-Smit, B. L., Wolters, E. C., and Berendse, H. W. (2004). Idiopathic hyposmia as a preclinical sign of Parkinson's disease. *Ann. Neurol.* 56, 173–181. doi: 10. 1002/ana. 20160

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Prins, D., Plank, T., Baseler, H. A., Gouws, A. D., Beer, A., Morland, A. B., et al. (2016). Surface-based analyses of anatomical properties of the visual cortex in macular degeneration. *PLoS One* 11: e0146684. doi: 10. 1371/journal. pone. 0146684

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Raz, N., Lindenberger, U., Rodriguez, K. M., Kennedy, K. M., Head, D., Williamson, A., et al. (2005). Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cereb. Cortex* 15, 1676–1689. doi: 10. 1093/cercor/bhi044

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Reyes-Ortiz, C. A., Kuo, Y. F., DiNuzzo, A. R., Ray, L. A., Raji, M. A., and Markides, K. S. (2005). Near vision impairment predicts cognitive decline: data from the Hispanic established populations for epidemiologic studies of the elderly. *J. Am. Geriatr. Soc.* 53, 681–686. doi: 10. 1111/j. 1532-5415. 2005. 53219. x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Roberts, R. O., Christianson, T., Kremers, W. K., Mielke, M. M., Machulda, M. M., Vassilaki, M., et al. (2016). Association between olfactory dysfunction and amnestic mild cognitive impairment and Alzheimer disease dementia. *JAMA Neurol.* 73, 93–101. doi: 10.1001/jamaneurol.2015.2952

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Rogers, M. A., and Langa, K. M. (2010). Untreated poor vision: a contributing factor to late-life dementia. *Am. J. Epidemiol.* 171, 728–735. doi: 10.1093/aje/kwp453

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Ross, G. W., Abbott, R. D., Petrovitch, H., Tanner, C. M., Davis, D. G., Nelson, J., et al. (2006). Association of olfactory dysfunction with incidental Lewy bodies. *Mov. Disord.* 21, 2062–2067. doi: 10.1002/mds.21076

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Saez de Asteasu, M. L., Martinez-Velilla, N., Zambom-Ferraresi, F., Casas-Herrero, A., and Izquierdo, M. (2017). Role of physical exercise on cognitive function in healthy older adults: a systematic review of randomized clinical trials. *Ageing Res. Rev.* 37, 117–134. doi: 10.1016/j.arr.2017.05.007

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Salvi, S. M., Akhtar, S., and Currie, Z. (2006). Ageing changes in the eye. *Postgrad. Med. J.* 82, 581–587. doi: 10.1136/pgmj.2005.040857

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Sapolsky, R. M., Krey, L. C., and McEwen, B. S. (1984). Stress down-regulates corticosterone receptors in a site-specific manner in the brain. *Endocrinology* 114, 287–292. doi: 10.1210/endo-114-1-287

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Saunders, G. H., and Echt, K. V. (2007). An overview of dual sensory impairment in older adults: perspectives for rehabilitation. *Trends Amplif.* 11, 243–258. doi: 10.1177/1084713807308365

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Scholey, A., Haskell, C., Robertson, B., Kennedy, D., Milne, A., and Wetherell, M. (2009). Chewing gum alleviates negative mood and reduces cortisol during acute laboratory psychological stress. *Physiol. Behav.* 97, 304–312. doi: 10.1016/j.physbeh.2009.02.028

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Schubert, C. R., Carmichael, L. L., Murphy, C., Klein, B. E. K., Klein, R., and Cruickshanks, K. J. (2008). Olfaction and the 5-year incidence of cognitive impairment in an epidemiologic study of older adults. *J. Am. Geriatrics Soc.* 56, 1517–1521. doi: 10.1111/j.1532-5415.2008.01826.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Schubert, C. R., Fischer, M. E., Pinto, A. A., Klein, B. E. K., Klein, R., and Cruickshanks, K. J. (2017). Odor detection thresholds in a population of older adults. *Laryngoscope* 127, 1257–1262. doi: 10.1002/lary.26457

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Segovia, G., Yague, A. G., Garcia-Verdugo, J. M., and Mora, F. (2006). Environmental enrichment promotes neurogenesis and changes the extracellular concentrations of glutamate and GABA in the hippocampus of aged rats. *Brain Res. Bull.* 70, 8–14. doi: 10.1016/j.brainresbull.2005.11.005

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Segura, B., Baggio, H. C., Solanaa, E., Palacios, E. M., Vendrell, P., Bargalló, N., et al. (2013). Neuroanatomical correlates of olfactory loss in normal aged subjects. *Behav. Brain Res.* 246, 148–153. doi: 10.1016/j.bbr.2013.02.025

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Seubert, J. J., Laukka, E. J., Rizzuto, D., Hummel, T., Fratiglioni, L., Bäckman, L., et al. (2017). Prevalence and correlates of olfactory dysfunction in old age: a population-based study. *J. Gerontol. A Biol. Sci. Med. Sci.* 72, 1072–1079. doi: 10.1093/gerona/glx054

[CrossRef Full Text](#) | [Google Scholar](#)

Shah, R. J., Diwan, F. J., Diwan, M. J., Chauhan, V. J., Agrawal, H. S., and Patel, G. C. (2015). A study of the emotional effects of tooth loss in an edentulous

Gujarati population and its association with depression. *J. Indian Prosthodont. Soc.* 15, 237–243. doi: 10. 4103/0972-4052. 161564

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Shen, J., Kassir, M. A., Wu, J., Zhang, Q., Zhou, S., Xuan, S. Y., et al. (2013). MR volumetric study of piriform-cortical amygdala and orbitofrontal cortices: the aging effect. *PLoS One* 8: 374526. doi: 10. 1371/journal. pone. 0074526

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Smith, A. (2010). Effects of chewing gum on cognitive function, mood and physiology in stressed and non-stressed volunteers. *Nutr. Neurosci.* 13, 7–16. doi: 10. 1179/147683010X12611460763526

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Sohrabi, H. R., Bates, K. A., Weinborn, M. G., Johnston, A. N. B., Bahramian, A., Taddei, K., et al. (2012). Olfactory discrimination predicts cognitive decline among community-dwelling older adults. *Transl. Psychiatry* 2: e118. doi: 10. 1038/tp. 2012. 43

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Sorrells, S. F., Paredes, M. F., Cebrian-Silla, A., Sandoval, K., Qi, D., Kelley, K. W., et al. (2018). Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555, 377–381. doi: 10. 1038/nature25975

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Spaccavento, S., Del Prete, M., Craca, A., and Fiore, P. (2009). Influence of nutritional status on cognitive, functional and neuropsychiatric deficits in Alzheimer's disease. *Arch. Gerontol. Geriatr.* 48, 356–360. doi: 10.1016/j.archger.2008.03.002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Spierer, O., Fischer, N., Barak, A., and Belkin, M. (2016). Correlation between vision and cognitive function in the elderly: a cross-sectional study. *Medicine* 95: e2423. doi: 10.1097/MD.0000000000002423

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Stathokostas, L., Theou, O., Little, R. M., Vandervoort, A. A., and Raina, P. (2013). Physical activity-related injuries in older adults: a scoping review. *Sports Med.* 43, 955–963. doi: 10.1007/s40279-013-0076-3

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Stein, P. S., Desrosiers, M., Donegan, S. J., Yepes, J. F., and Kryscio, R. J. (2007). Tooth loss, dementia and neuropathology in the Nun study. *J. Am. Dent. Assoc.* 138, 1314–1322. doi: 10.14219/jada.archive.2007.0046

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Stephens, R., and Tunney, R. J. (2004). Role of glucose in chewing gum-related facilitation of cognitive function. *Appetite* 43, 211–213. doi: 10.1016/j.appet.2004.07.006

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Stewart, R., Stenman, U., Hakeberg, M., Hägglin, C., Gustafson, D., and Skoog, I. (2015). Associations between oral health and risk of dementia in a 37-year follow-up study: the prospective population study of women in Gothenburg. *J. Am. Geriatr. Soc.* 63, 100–105. doi: 10.1111/jgs.13194

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Stimpson, N. J., Davison, G., and Javadi, A. H. (2018). Jogging' the noggin: towards a physiological understanding of exercise-induced cognitive benefits. *Neurosci. Biobehav. Rev.* 88, 177–186. doi: 10.1016/j.neubiorev.2018.03.018

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Sungkarat, S., Boripuntakul, S., Kumfu, S., Lord, S. R., and Chattipakorn, N. (2018). Tai chi improves cognition and plasma BDNF in older adults with mild cognitive impairment: a randomized controlled trial. *Neurorehabil. Neural Repair.* 32, 142–149. doi: 10.1177/1545968317753682

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Syka, J. (2002). Plastic changes in the central auditory system after hearing loss, restoration of function and during learning. *Physiol. Rev.* 82, 601–636. doi: 10.1152/physrev.00002.2002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Syrjälä, A. M., Ylöstalo, P., Sulkava, R., and Knuutila, M. (2007). Relationship between cognitive impairment and oral health: results of the Health 2000

Health Examination Survey in Finland. *Acta Odontol. Scand.* 65, 103–108.  
doi: 10.1080/00016350601083521

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Tabaton, M., Monaco, S., Cordone, M. P., Colucci, M., Giaccone, G., Tagliavini, F., et al. (2004). Prion deposition in olfactory biopsy of sporadic Creutzfeldt-Jakob disease. *Ann. Neurol.* 55, 294–296. doi: 10.1002/ana.20038

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Takeda, Y., Oue, H., Okada, S., Kawano, A., Koretake, K., Michikawa, M., et al. (2016). Molar loss and powder diet leads to memory deficit and modifies the mRNA expression of brain-derived neurotrophic factor in the hippocampus of adult mice. *BMC Neurosci.* 17: 81. doi: 10.1186/s12868-016-0319-y

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Tamura, H., Tsukamoto, H., Mukai, S., Kato, T., Minamoto, A., Ohno, Y., et al. (2004). Improvement in cognitive impairment after cataract surgery in elderly patients. *J. Cataract. Refract. Surg.* 30, 598–602. doi: 10.1016/j.jcrs.2003.10.019

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Tesch-Römer, C. (1997). Psychological effects of hearing aid use in older adults. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 52, 127–138. doi: 10.1093/geronb/52B.3.P127

[CrossRef Full Text](#) | [Google Scholar](#)

Thomson, R. S., Aduong, P., Miller, A. T., and Gurgel, R. K. (2017). Hearing loss as a risk factor for dementia: a systematic review. *Laryngoscope Investig. Otolaryngol.* 2, 69–79. doi: 10.1002/lio2.65

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Tonetti, M. S., Jepsen, S., Jin, L., and Otomo-Corgel, J. (2017). Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: a call for global action. *J. Clin. Periodontol.* 44, 456–462. doi: 10.1111/jcpe.12732

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Toussaint, N., de Roon, M., van Campen, J. P., Kremer, S., and Boesveldt, S. (2015). Loss of olfactory function and nutritional status in vital older adults and geriatric patients. *Chem. Senses* 40, 197–203. doi: 10.1093/chemse/bju113

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Tucha, O., Mecklinger, L., Maier, K., Hammerl, M., and Lange, K. (2004). Chewing gum differentially affects aspects of attention in healthy subjects. *Appetite* 42, 327–329. doi: 10.1016/j.appet.2004.01.003

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Uhlmann, R. F., Larson, E. B., Koepsell, T. D., Rees, T. S., and Duckert, L. G. (1991). Visual impairment and cognitive dysfunction in Alzheimer's disease. *J. Gen. Intern. Med.* 6, 126–132. doi: 10.1007/BF02598307

[CrossRef Full Text](#) | [Google Scholar](#)

Utsugi, C., Miyazono, S., Osada, K., Sasajima, H., Noguchi, T., Matsuda, M., et al. (2014). Hard-diet feeding recovers neurogenesis in the subventricular zone and olfactory functions of mice impaired by soft-diet feeding. *PLoS One* 9: e97309. doi: 10.1371/journal.pone.0097309

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Vaden, K. I., Kuchinsky, S. E., Ahlstrom, J. B., Teubner-Rhodes, S. E., Dubno, J. R., and Eckert, M. A. (2016). Cingulo-opercular function during word recognition in noise for older adults with hearing loss. *Exp. Aging Res.* 42, 86–106. doi: 10.1080/0361073X.2016.1108784

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Vaden, K. I., Kuchinsky, S. E., Cute, S. L., Ahlstrom, J. B., Dubno, J. R., and Eckert, M. A. (2013). The cingulo-opercular network provides word-recognition benefit. *J. Neurosci.* 33, 18979–18986. doi: 10.1523/JNEUROSCI.1417-13.2013

[CrossRef Full Text](#) | [Google Scholar](#)

van Hooren, S. A., Anteunis, L. J., Valentijn, S. A., Bosma, H., Ponds, R. W., Jolles, J., et al. (2005). Does cognitive function in older adults with hearing impairment improve by hearing aid use? *Int. J. Audiol.* 44, 265–271. doi: 10.1080/14992020500060370

[CrossRef Full Text](#) | [Google Scholar](#)

Varin, M., Kergoat, M. J., Belleville, S., Li, G., Rousseau, J., Roy-Gagnon, M. H., et al. (2017). Age-related eye disease and participation in cognitive activities. *Sci. Rep.* 7: 17980. doi: 10.1038/s41598-017-18419-2

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Vaynman, S., Ying, Z., and Gómez-Pinilla, F. (2004). Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur. J. Neurosci.* 20, 2580–2590. doi: 10.1111/j.1460-9568.2004.03720.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Voss, M. W., Prakash, R. S., Erickson, K. I., Basak, C., Chaddock, L., Kim, J. S., et al. (2010). Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Front. Aging Neurosci.* 26: 32. doi: 10.3389/fnagi.2010.00032

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Walls, A. W., Steele, J. G., Sheiham, A., Marcene, W., and Moynihan, P. J. (2000). Oral health and nutrition in older people. *J. Public Health Dent.* 60, 304–307. doi: 10.1111/j.1752-7325.2000.tb03339.x

[CrossRef Full Text](#) | [Google Scholar](#)

Wattendorf, E., Welge-Lussen, A., Fiedler, K., Bilecen, D., Wolfensberger, M., Fuhr, P., et al. (2009). Olfactory impairment predicts brain atrophy in Parkinson's disease. *J. Neurosci.* 29, 15410–15413. doi: 10.1523/JNEUROSCI.1909-09.2009

[CrossRef Full Text](#) | [Google Scholar](#)

Wei, J., Hu, Y., Zhang, L., Hao, Q., Yang, R., Lu, H., et al. (2017). Hearing impairment, mild cognitive impairment, and dementia: a meta-analysis of cohort studies. *Dement. Geriatr. Cogn. Dis. Extra* 7, 440–452. doi: 10.1159/000485178

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Weijenberg, R. A., Scherder, E. J., and Lobbezoo, F. (2011). Mastication for the mind - the relationship between mastication and cognition in ageing and dementia. *Neurosci. Biobehav. Rev.* 35, 483–497. doi: 10.1016/j.neubiorev.2010.06.002

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Welmer, A. K., Rizzuto, D., Parker, M. G., and Xu, W. (2017). Impact of tooth loss on walking speed decline over time in older adults: a population-based cohort study. *Aging Clin. Exp. Res.* 29, 793–800. doi: 10.1007/s40520-016-0630-6

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

White-Schwoch, T., Carr, K. W., Anderson, S., Strait, D. L., and Kraus, N. (2013). Older adults benefit from music training early in life: biological evidence for long-term training-driven plasticity. *J. Neurosci.* 33, 17667–17674. doi: 10.1523/JNEUROSCI.2560-13.2013

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Whitson, H. E., Ansah, D., Whitaker, D., Potter, G., Cousins, S. W., MacDonald, H., et al. (2010). Prevalence and patterns of comorbid cognitive impairment in low vision rehabilitation for macular disease. *Arch. Gerontol. Geriatr.* 50, 209–212. doi: 10.1016/j.archger.2009.03.010

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Whitson, H. E., Whitaker, D., Sanders, L. L., Potter, G. G., Cousins, S. W., Ansah, D., et al. (2012). Memory deficit is associated with worse functional trajectories among older adults in low vision rehabilitation for macular disease. *J. Am. Geriatr. Soc.* 60, 2087–2092. doi: 10.1111/j.1532-5415.2012.04194.x

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

WHO (2018). *Deafness and Hearing Loss Fact Sheet*. Geneva: WHO.

[Google Scholar](#)

Williamson, R. A., Pytynia, K., Oghalai, J. S., and Vrabec, J. T. (2009). Auditory performance after cochlear implantation in late septuagenarians and octogenarians. *Otol. Neurotol.* 30, 916–920. doi: 10.1097/MAO.0b013e3181b4e594

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Wong, P. C. M., Ettlinger, M., Sheppard, J. P., Gunasekera, G. M., and Dhar, S. (2010). Neuroanatomical characteristics and speech perception in noise in older adults. *Ear Hear.* 31, 471–479. doi: 10.1097/AUD.0b013e3181d709c2

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Woo, C. C., Donnelly, J. H., Steinberg-Epstein, R., and Leon, M. (2015). Environmental enrichment as a therapy for autism: a clinical trial replication and extension. *Behav. Neurosci.* 129, 412–422. doi: 10.1037/bne0000068

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Woo, C. C., and Leon, M. (2013). Environmental enrichment as an effective treatment for autism: a randomized controlled trial. *Behav. Neurosci.* 127, 487–497. doi: 10.1037/a0033010

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Yamamoto, T., and Hirayama, A. (2001). Effects of soft-diet feeding on synaptic density in the hippocampus and parietal cortex of senescence-accelerated mice. *Brain Res.* 902, 255–263. doi: 10.1016/S0006-8993(01)02410-6

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Yao, L., Pinto, J. M., Xiaoli, Y., Li, L., Peng, P., and Wei, Y. (2014). Gray matter volume reduction of olfactory cortices in patients with idiopathic olfactory loss. *Chem. Senses* 39, 755–760. doi: 10.1093/chemse/bju047

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Yin, Y., Hou, Z., Wang, X., Sui, Y., and Yuan, Y. (2015). Association between altered resting-state cortico-cerebellar functional connectivity networks and

mood/cognition dysfunction in late-onset depression. *J. Neural Transm.* 122, 887–896. doi: 10.1007/s00702-014-1347-3

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Young, J., Angevaren, M., Rusted, J., and Tabet, N. (2015). Aerobic exercise to improve cognitive function in older people without known cognitive impairment. *Cochrane Database Syst. Rev.* 4: CD005381. doi: 10.1002/14651858.CD005381.pub4

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Yu, H., Chen, X., Liu, J., and Zhou, X. (2013). Gum chewing inhibits the sensory processing and the propagation of stress-related information in a brain network. *PLoS One* 8: e57111. doi: 10.1371/journal.pone.0057111

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Yuan, J., Sun, Y., Sang, S., Pham, J. H., and Kong, W. J. (2018). The risk of cognitive impairment associated with hearing function in older adults: a pooled analysis of data from eleven studies. *Sci. Rep.* 8: 2137. doi: 10.1038/s41598-018-20496-w

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Zendel, B. R., and Alain, C. (2012). Musicians experience less age-related decline in central auditory processing. *Psychol. Aging* 27, 410–417. doi: 10.1037/a0024816

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Zheng, Y., Fan, S., Liao, W., Fang, W., Xiao, S., and Liu, J. (2017). Hearing impairment and risk of Alzheimer's disease: a meta-analysis of prospective cohort studies. *Neurol. Sci.* 38, 233–239. doi: 10.1007/s10072-016-2779-3

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)

Zhou, Q., and Faure Walker, N. (2015). Promoting vision and hearing aids use in an intensive care unit. *BMJ Qual. Improv. Rep.* 4: u206276. w2702. doi: 10.1136/bmjquality. u206276. w2702

[PubMed Abstract](#) | [CrossRef Full Text](#) | [Google Scholar](#)